

UNIVERSITY
OF MICHIGAN

MAY 25 1953

ASTRONOMY
LIBRARY

Sky and **TELESCOPE**



A family of telescopes

In This Issue:

★
Vol. XII, No. 8

JUNE, 1953

★
Whole Number 140

The Distance Scale
of the Universe — I

Wilkie Observatory

Student Observations
of Delta Cephei

Stones from the Sky?

Next Year's Favorable
Total Eclipse

Northern and Southern
Star Charts

NEW PRECISION OSCILLATOR

Eclipse observers, usually in need of high-precision clocks, will welcome a new portable continuously oscillating frequency standard recently developed by Peter G. Sulzer, of the National Bureau of Standards. With accuracy it combines small size, low weight, and easy temperature control.

Fitting into a tube less than two inches in diameter and seven inches long, the major components of the new transistor oscillator are a type 2517 junction transistor, a high-precision 100-kilocycle GT-cut quartz crystal unit, and a long-life mercury cell. Operating at 1.35 volts at 100 microamperes, the dry-cell power supply should have an active life of five or more years.

Over half of the space in the tube is occupied by the crystal, which is mounted in an evacuated glass envelope. The transistor, coil, capacitors, and resistors are supported on a bakelite frame that may be "potted" in casting resin to add rigidity to the section. The mercury cell, only about one-half inch deep, is at the base of the assembly and is insulated from the metal "can" by a bakelite shield. A constant phase shift in the feedback loop associated with the crystal is obtained by using large, stable "swamping" capacitors at both crystal connections and by means of highly stable components in the remainder of the circuit. Constancy of amplitude of oscillation is achieved by operating the transistor in such a manner that collector-voltage limiting is produced. The diagram shows the manner in which the 100-kilocycle tank circuit voltage is reduced by means of an attenuator.

Over short intervals, time variations of about three parts in 10^{10} have been found in the oscillator, and the drift in 24 hours is about three parts in 10^9 . These figures are comparable to those obtained from vacuum-tube standard oscillators, particularly at the time of their initial installation; frequency drift normally decreases with age. The change in frequency is about one part in 10^8 per 0.10-volt variation in the supply voltage.


In the field it is always difficult to control the temperature of equipment. Standard quartz oscillators and quartz clocks have heretofore required relatively complex temperature control apparatus (operating at temperatures up to 60° centigrade). With a change in temperature of one degree, the new oscillator's frequency may vary one part in 10⁸, but reasonable stability can be achieved by merely placing the oscillator in a Dewar flask containing crushed clear ice (0° centigrade). Thus, it now becomes possible to carry this frequency standard anywhere in the world. D.H.

Sky and TELESCOPE

Copyright, 1953, by
SKY PUBLISHING CORPORATION

CHARLES A. FEDERER, JR., *Editor*; HELEN S. FEDERER, *Managing Editor*

EDITORIAL ADVISORY BOARD: *Clement S. Brainin*, Amateur Astronomers Association, New York; *Edward A. Halbach*, Milwaukee Astronomical Society; *Donald H. Menzel*, Harvard College Observatory; *Paul W. Merrill*, Mount Wilson Observatory (retired); *Charles H. Smiley*, Ladd Observatory; *Percy W. Witherell*, Bond Astronomical Club.



Components and circuit of National Bureau of Standards transistor oscillator.

VOL. XII, No. 8
WHOLE NUMBER 140

CONTENTS

JUNE, 1953

COVER: Four portable reflecting telescopes built by James V. Lawrence, Flushing, N. Y., whose mountings, consisting principally of masonite, permit observing from the southern horizon to the zenith when operated as equatorials. The mirrors are 3-inch, 4-inch, 5-inch, and 6-inch, respectively. (See page 219.)

THE DISTANCE SCALE OF THE UNIVERSE — I — Otto Struve	203
WILKIE OBSERVATORY — D. C. Dornberg	207
STUDENT OBSERVATIONS OF DELTA CEPHEI — Peter van de Kamp	208
STONES FROM THE SKY? — Helmut Hecksher	210
NEXT YEAR'S FAVORABLE TOTAL ECLIPSE — Paul W. Stevens	214
Amateur Astronomers	213
Books and the Sky	216
Gleanings for ATM's	219
Stars for June	227
News Notes	206
Observer's Page	223
Southern Stars	226

BACK COVER: The total solar eclipse of February 25, 1952, photographed at Khartoum, Anglo-Egyptian Sudan, by Dr. G. Van Biesbroeck, of Yerkes Observatory, University of Chicago, for the purpose of checking the Einstein shift of starlight passing the sun. The plate was exposed 1½ minutes through a yellow filter on Eastman 103a-E emulsion, with maximum sensitivity around 6500 angstroms, for which wave length the lens was computed. The print was made with a specially cut rotating sector to procure proper rendition of both the bright inner regions and the faint outer extensions of the corona, which required a difference of 100 in exposure time. Note the fine structure of the fan-shaped polar spikes, and the very long equatorial streamers typical at sunspot minimum. Photograph, courtesy National Geographic Society.

SKY AND TELESCOPE is published monthly by Sky Publishing Corporation, Harvard College Observatory, Cambridge 38, Mass. Entered as second class matter, April 23, 1939, at the Post Office, Boston, Mass., under Act of March 3, 1879; accepted for mailing at the special rate of postage provided in Paragraph 4, Section 538, Postal Laws and Regulations.

Subscriptions: \$4.00 per year in the United States and possessions, and to Latin-American countries; \$7.00 for two years. Add \$1.00 per year for Canada and for all other foreign countries, making the total subscription \$5.00 per year and \$9.00 for two years. Canadian and foreign remittances should be made in United States currency. Single copies, 35 cents, foreign 45 cents. Circulation staff: Betty G. Dodd, manager; Nancy R. Bolton; Virginia K. Cox.

Betty G. Dodd, manager; Nancy R. Bolton; Virginia K. Cox.
Editorial and advertising offices: Harvard College Observatory, Cambridge 38, Mass. Unsolicited articles and pictures are welcome, bearing adequate return postage, but we cannot guarantee prompt editorial attention, nor are we responsible for the return of unsolicited manuscripts.
The articles in SKY and TELESCOPE, beginning with Vol. XII, are indexed in THE READERS' GUIDE TO PERIODICAL LITERATURE.

THE YEAR 1952 will go down in astronomical history as one in which astronomers accepted a tremendous change in their concept of the distance scale of the universe. Also, it will mark an astronomical milestone that we have reached directly as a result of observations with the 200-inch Hale telescope — observations which no other instrument in the world is capable of making. The significance of this change is concisely described in the following paragraph quoted directly from the official minutes of Commission 28, Extragalactic Nebulae, of the International Astronomical Union, at its Rome meeting, September 4-13, 1952:

"Dr. Baade then went on to describe several results of great cosmological significance. He pointed out that, in the course of his work on the two stellar populations in M31, it had become more and more clear that either the zero point of the classical Cepheids or the zero point of the cluster variables must be in error. Data obtained recently — Sandage's color-magnitude diagram of M3 — supported the view that the error lay with the zero point of the classical Cepheids, not with the cluster variables. Moreover, the error must be such that our previous estimates of extragalactic distances — not distances within our own galaxy — were too small by as much as a factor 2. Many notable implications followed immediately from the corrected distances: the globular clusters in M31 and in our own galaxy now come out to have closely similar luminosities; and our galaxy may now come out to be somewhat smaller than M31. Above all, Hubble's characteristic time-scale for the universe must now be increased from about 1.8×10^9 years to about 3.6×10^9 years."

In the final analysis, our knowledge of the distances of stars and galaxies rests upon the determination of the scale of the solar system; for several hundred years astronomers have been concerned with this first stage of the problem, until now we know solar system distances with good accuracy. Then, using as a base line the astronomical unit, that is, the distance between the earth and the sun, astronomers succeeded in 1837 in measuring the parallax displacement and, hence, the distance of one of the nearest of the stars. The trigonometric parallax determinations which followed in the next 116 years form the second stage in the exploration of the universe.

The third stage resulted from Miss H. Leavitt's discovery, in 1912, of a close relation between the apparent brightnesses and the periods of Cepheid variable stars in the Small Magellanic Cloud. From a study of 25 variables, she made a diagram in which the apparent magnitude of each star at maximum and at minimum light was plotted against the logarithm of the period. A



The new distance scale of the universe places the Great Nebula in Andromeda at a distance of about 1,500,000 light-years, and increases its known size to somewhat more than that of our own Milky Way system. This photograph of M31 and its companion galaxies was taken with the Jewett Schmidt telescope of Harvard Observatory.

THE DISTANCE SCALE OF THE UNIVERSE -- I

BY OTTO STRUVE, *Leuschner Observatory*
University of California


variable with a period of exactly 10 days would have a logarithm of the period equal to 1.0 and would, according to the diagram, appear on a photographic plate with an apparent magnitude of 13.5 when at maximum light and about 14.7 when at minimum light. Fainter variables were found to have shorter periods;

those of magnitudes between 15 at maximum and about 16.5 at minimum correspond to a period of about one day.

This diagram was the beginning of all our recent knowledge concerning the distances between our galaxy and other galaxies such as the Andromeda nebula, M31, and the great spiral in Triangulum, M33. Although at the time Miss Leavitt made this discovery, the importance of the period-luminosity relation for distance measurements was not appreciated, either by her or by E. C. Pickering, then the director of the Harvard Observatory, Miss Leavitt did remark:

"Since the variables are probably at nearly the same distance from the earth, their periods are apparently associated with their actual emission of light, as determined by their mass, density, and surface brightness. . . Two fundamental questions upon which light may be thrown . . . are whether there are definite limits to the mass of variable stars of the cluster type, and if the spectra of such variables having long periods differ from those . . . whose periods are short."

It should be realized that Miss Leavitt's diagram as such did not directly



The original period-luminosity curve, copied from Harvard Observatory "Circular" No. 173, March 3, 1912, in which Miss Leavitt's discoveries are described.



The Small Magellanic Cloud, in which are contained the variable stars used by Miss Leavitt to discover the period-luminosity relation. Harvard Observatory photograph.

provide the means for determining distances. It merely showed, for instance, that a Cepheid whose period is 10 times longer than that of another variable of this kind is approximately two magnitudes brighter than the latter. Two magnitudes correspond to a difference in intrinsic luminosity of $6\frac{1}{4}$ times. Thus, if we observe two variable stars of the same apparent magnitude at maximum light, the one having a period of four days, the other of 40 days, we can conclude that the longer-period variable is $6\frac{1}{4}$ times as bright but considerably farther away. As brightness varies inversely with the square of the distance, the actual relative distance would be the square root of $6\frac{1}{4}$, or about $2\frac{1}{2}$ times; the star of longer period must be $2\frac{1}{2}$ times as distant as the one of shorter period.

But if we have not yet determined the distance of the nearer star, our observation provides only relative distances and does not give us any information regarding the absolute scale of distances. This latter and important aspect of the period-luminosity relation of the Cepheid variables is usually described as setting the zero point of the period-luminosity curve.

The extension of Miss Leavitt's work and its practical use as a yardstick for measuring the distances of globular clusters and extragalactic systems was made by H. Shapley in a long series of papers

published in the *Astrophysical Journal* more than 30 years ago, while he was a member of the Mount Wilson Observatory staff. In some of his early work he determined the zero point, although this had also, in effect, been done by E. Hertzsprung in 1913 on the basis of the Boss proper motions for typical Cepheids.


If we could measure trigonometrically the distance of a single Cepheid variable star in our Milky Way system, and if we could estimate the amount of interstellar absorption between us and this variable

star, we would have an immediate determination of the zero point. Unfortunately, even the brightest of the Cepheid variables are so far away that ordinary trigonometric parallaxes are useless. Nor is there any indirect method available of the kind astronomers use when they take advantage of a special type of star occurring as a member of a visual binary system. Cepheid variables do not even occur among moving clusters of stars, such as the Hyades, or among other galactic clusters, like the Pleiades, for which the distances can be estimated with the help of the Hertzsprung-Russell diagram (*Sky and Telescope*, May, 1951, page 164). We are left with the difficult method of estimating statistically the distances of certain groups of galactic Cepheids with the help of their radial velocities and proper motions.

The method is, in principle, quite simple. Suppose you are located on a high building and you watch the movement of persons in the streets below. Suppose you know that the average speed of a person walking is one yard per second; you will also know that some persons are walking considerably faster than the average and others much slower. Therefore, you observe some of them to change their positions hardly at all, while others appear displaced about 20 degrees in 10 seconds. Between these extremes there are individuals that appear to move with various speeds, and you conclude that their average apparent displacement is 10 degrees in 10 seconds or one degree in one second. You would further conclude that, on the average, a displacement of one degree per second corresponds to the known average linear velocity of one yard per second. In other words, at your distance above the street, one yard subtends an angle of one degree; from this you can easily compute that your elevation is about 60 yards.

The apparent angular motions of the bright Cepheid variables have been determined from accurate measures of


The photographic period-luminosity relation adopted by Harlow Shapley in his book, "Star Clusters," 1930. The dots represent means for intervals of logarithms of the periods; the crosses, for intervals of absolute magnitude.



their positions in the sky extending over hundreds of years. For the same stars, motions in the line of sight, or radial velocities, have been determined from measures of the Doppler shifts of their spectral lines. An individual Cepheid, however, may be moving only at right angles to the line of vision, thus having large proper motion and no radial velocity with respect to the earth. Another star could be moving entirely along the line of sight, and would then have no detectable proper motion or tangential velocity. For an individual star we cannot determine whether a small proper motion results from an actual small velocity at right angles to the line of sight, or whether the motion seems small because the star is very far away. But if we consider the problem statistically, we may assume the motions to be at random, as large a component being radial as tangential, and we may roughly equate the radial and linear tangential velocities. On this basis, the measured radial velocity should be approximately a measure of the linear motion of the star, while the angular proper motion should be a measure of this same velocity as seen from the distance between the earth and the star. Thus, if a star of large radial velocity has a small proper motion, its distance must be relatively great; stars of larger proper motion must be nearer.

The mathematics of this process is not difficult to understand, but it is unnecessary for our present purpose. The important point is to realize that if, over a large statistical sample of stellar data, we grant the similarity of the radial and tangential velocities, relatively reliable statistical parallaxes may be obtained.

In this manner, Shapley set the zero point of his famous period-luminosity curve, in which the abscissa has the same meaning as in the period-luminosity relation of Miss Leavitt, but the ordinate now represents the intrinsic or absolute photographic magnitudes of the variable stars half way between maximum and minimum light. This diagram, therefore, gave us a means of determining the individual distances of the Cepheid variable stars once we observed their periods.



Here is shown the relation among cross motion (tangential velocity), radial velocity, and the space motion of a star. Proper motion is the angular change of position produced by the cross motion, but its observed value depends also on the distance of the star.



The Large Magellanic Cloud, pictured here, and the Small Cloud are now considered to be at a distance of about 150,000 light-years, or 1/10 as far from the earth as the Great Nebula in Andromeda. Harvard Observatory photograph.

Let us suppose that in our galaxy we have observed a variable star whose apparent magnitude, half way between maximum and minimum light, is $+8.2$. After a series of observations we find that the period is 10 days. Taking the logarithm of the period, namely 1.0, and entering with it Shapley's diagram, we read off the ordinate scale that the absolute magnitude of the variable is -1.8 . The star thus appears 10 magnitudes fainter when actually observed on a photographic plate than it would be if it were located at the standard absolute magnitude distance of 10 parsecs or 33 light-years. A difference of 10 in magnitude corresponds to a ratio in light of 10,000. Since light intensity diminishes with the square of the distance, the real distance of this variable is 100 times greater than 10 parsecs and is thus equal to 1,000 parsecs.

Since we know that interstellar absorption tends to make the stars look fainter than they would otherwise, we have to introduce a correction to the observed apparent magnitude, allowing for intervening dust and gas, and thus reducing the distance somewhat.

This method is perfectly straightforward, and it was used successfully at Mount Wilson by E. P. Hubble and all of his followers in determining the distances of the nearer galaxies. In these systems many Cepheid variables were discovered, their periods were deter-

mined, and the rest of the procedure resembled the example we have worked out in the preceding paragraphs.

In this manner Hubble, in 1929, found that Cepheids of corresponding period in the Andromeda galaxy are about 4.6 magnitudes fainter than in the Small Magellanic Cloud. This would make the distance of the Andromeda galaxy $8\frac{1}{2}$ times greater than the distance of the cloud. Using Shapley's value of 106,000 light-years for the cloud, Hubble set the distance of the Andromeda galaxy as 900,000 light-years.

Except for a relatively minor correction in this figure—bringing it down to 750,000 light-years—which is due to the small amount of interstellar absorption in the direction of Andromeda, this result remained generally accepted until Baade's report at the Rome meeting. Not only that, but all other extragalactic distances were tied to the distance found by Hubble with the help of the Cepheid variables. Hence, the framework of the metagalactic system rested entirely upon Shapley's zero point of the period-luminosity law.

Next month we shall see how the necessity to change the zero point had become increasingly apparent, and how the 200-inch telescope demonstrated the need to change our concept of the size of the universe by a factor of two.

(To be concluded)

NEWS NOTES

LUNAR PROFILES FROM A BEADED ECLIPSE

The solar eclipse of May 9, 1948, was annular, with 0.9996 of the sun's diameter eclipsed as seen from Rebun Island, Hokkaido, Japan. Thus at maximum phase little more of the sun could be seen than Baily's beads around the whole perimeter. Over 200 separate beads were observed. In the *Publications* of the Astronomical Society of Japan, Shigetsugu Fujinami, of Kyoto Observatory, describes how times of contacts at this eclipse were utilized for the determination of the heights of some of the mountains and depths of deep valleys at the limb of the moon.

Moving pictures at the rate of 24 frames a second for an interval of one minute bracketing the maximum phase were taken with a Cassegrain reflector with a focal length of 1,400 millimeters. For the first and last contacts the exposures were $f/18$, $1/432$ second, and during the beaded phases $f/15$, $1/50$ second. Time markers were printed along the film sound track at half-second intervals to an accuracy better than 0.02 second. The times of contact could readily be reduced to heights above a mean lunar limb. Such contour data have ordinarily been obtained from measurements of photographs of the moon, and standard results are available in Friedrich Hayn's *Selenographische Koordinaten* (1914).

Fujinami finds that many of the mountains are appreciably higher and the valleys deeper than the older tables indicate. Moreover, analyzing the relative accuracies of the methods for determining the jagged lunar profiles, he is convinced that the eclipse method is substantially more accurate. Hayn's data are given to the nearest $1/10$ second of arc and Fujinami's to hundredths. Numerous conspicuous features

observed by both differ by 0.5 second or more. The most outstanding discrepancies found are for the depressions d'Alembert, Rook, and Mare Humboldtianum, respectively $2''.48$, $2''.34$, and $1''.29$ deeper than previously thought, and the mountain d'Alembert, which is $0''.68$ higher. Irradiation on photographs would appear to account for a smaller range of altitudes than that obtained from accurate time determinations. It will be of interest to see how Fujinami's results compare with photoelectrically determined profiles being obtained by C. B. Watts at the U. S. Naval Observatory.

IRISH SOLAR ECLIPSES

For the benefit of those concerned with chronology, F. J. O'Connor, of the Dunsink Observatory, has compiled in the *Proceedings* of the Royal Irish Academy a list of solar eclipses visible from Ireland between A.D. 400 and 1000. During this interval, there occurred a total of 1,421 eclipses of the sun visible from anywhere on the earth. Of these, 862 could immediately be eliminated as altogether invisible from Ireland.

For the remaining 559, the maximum phase was visible from Ireland in 165 cases, and in 28 more cases the sun could be seen after sunrise or before sunset at least 0.2 eclipsed. Thus, from an area of about 32,000 square miles, portions of 193 solar eclipses could definitely be seen out of a total of 1,421 in 600 years.

AWARD TO BART J. BOK

For a paper entitled, "Studies of the Southern Milky Way," Dr. Bart J. Bok, of Harvard Observatory, has been awarded the Lewis prize of the American Philosophical Society. Read at the society meeting on November 9, 1951, the paper deals with the equipment of the

Boyden station at Bloemfontein, South Africa, the major programs of the work then in progress, and the results of Dr. Bok's $1\frac{1}{2}$ years sojourn at that observatory. Analysis of the data that were accumulated during this relatively short period, when Dr. Bok's major function was to install and test the Armagh-Dunsink-Harvard telescope, will occupy his assistants for several years.

CHLORINE IN METEORITES

Several years ago, A. Gatterer and V. Frodl reported in *Ricerche Spettroscopiche* (a Vatican publication) on a spectroscopic technique for analyzing very minute amounts of nonmetallic substances by means of an electrodeless high-frequency discharge method.

In a current number of the same journal, E. Salpeter, of Cornell University, describes the application of this technique to the determination of the amount of chlorine in stony meteorites. He finds it eminently satisfactory: "... especially suitable for routine analyses and for cases when only a small amount of material is available." Twenty meteorites examined gave an average chlorine content of 0.09 per cent to an accuracy of 10 per cent. This value is in good agreement with previous determinations for stony meteorites.


CHLOROPHYLL FORMATION

The chemical basis to most forms of life, chlorophyll, could have been spontaneously created in the lifeless world of two billion or more years ago, according to Science Service, reporting the results of experiments at Ohio State University under the direction of Dr. William M. MacNevin. When carbon dioxide, ammonia, and water were passed over a heated silica tube (representing the hot rocks of the earth's early days), molecules of porphyrin were produced; this substance has a basic structure like that of chlorophyll.

In another experiment re-creating conditions in the earth's youth, artificial lightning was sent through an atmosphere of marsh gas (methane) and water vapor. It is believed that lightning was almost continuous in the atmosphere of the cooling earth. The experiment resulted in the formation of a resinous substance so complex chemically that its structure could not be analyzed.

NATURAL NEPTUNIUM 237

In 1947, Nobel prizewinner Glenn T. Seaborg predicted that the isotope of neptunium of atomic weight 237, first discovered with the Berkeley cyclotron in 1942, would be found in nature in minute quantities. This prediction has now been fulfilled by its discovery in ores in the Belgian Congo.



The middle of the annular eclipse of May 9, 1948, as it was photographed by a Kyoto Observatory expedition to Rebun Island. The limb features named are those for which large discrepancies in height or depth are found when compared with Hayn's data.

Wilkie Observatory

By D. C. DORNBERG

A NEW OBSERVATORY, housing an 8-inch reflecting telescope, has been added to Macalester College, St. Paul, Minn. The cost of the instruments and dome, with their operating mechanisms, was borne by the Wilkie Foundation. The Continental Machine Company, of Savage, Minn., one of the sponsors of the foundation, built the main telescope, the optics of which are by R. E. English, of St. Paul. Macalester College constructed the building.

Dr. W. J. Luyten, professor of astronomy at the University of Minnesota, is teaching the first course in astronomy. The Wilkie Observatory is under the direction of Dr. Waldo Glock, chairman of the department of geology and astronomy at Macalester.

Two consultants for maintenance and future development of the observatory were appointed by the college, Mr. English and the writer, who will teach an evening class in popular astronomy at the college. We are members of the St. Paul Telescope Club, as is Dr. Glock.

The ground floor of the new building is pear shaped and measures approximately 16 by 32 feet, the small end pointing east. The 12-foot dome rests on 8-inch reinforced concrete slab supported by crossed 8-inch I-beams.

Dr. Waldo Glock, chairman of the department of geology and astronomy at the college, is here adjusting the right-ascension drive of the 8-inch reflector.

Five smaller telescopes of 60 power each can readily be fastened to the observation platform railing east of the dome. These are used for student instruction, permitting a number to observe at one time.

The main telescope is an 8-inch Newtonian-Springfield of 64-inch focal length; it is equipped with a 20-power finder. The telescope pier is made of 8-inch welded steel pipe and rests on the intersection of the I-beams. Spindles $2\frac{1}{2}$ inches in diameter support both the right-ascension and declination axes,



serving as the bearings. The spindles were made a few 10-thousandths of an inch under size and were then chrome plated and lapped in. Thus, the fit allowance of the spindles is less than $5/10,000$ of an inch.

The setting circles are machine-indexed 10-inch disks of chrome-plated steel. A $1/20$ horsepower synchronous motor turns the right-ascension movement once in 24 hours. Thus, the telescope tracks at a mean solar instead of a sidereal rate, which is satisfactory for an instrument used primarily for instruction.

SHORT-PERIOD VISUAL BINARY

By means of an interferometer attached to the eyepiece of the $26\frac{1}{2}$ -inch refractor of the Union Observatory at Johannesburg, South Africa, W. S. Finsen has been conducting a survey for visual double stars so close together that they appear as one except when resolved into two stars by the interferometer technique. In the course of his survey, Dr. Finsen has now found the star Epsilon Ceti to be double, and in the February number of the *Monthly Notes* of the Astronomical Society of South Africa he gives a provisional orbit.

The period indicated, 1.59 years, is shorter than that of any other visual binary except Capella (an interferometric binary which has a period of only 105 days). The next shortest visual binary period is 1.715 years, determined by G. P. Kuiper for the star BD -8°-4352. Dr. Finsen is cautious in announcing his results, for they depend on only 21 observations spanning an interval of only slightly more than two years; some ambiguity in the interpretation of some of the measured position angles is possible. The star is very deserving of further investigation.



The Wilkie Observatory at Macalester College, which was dedicated in October, 1952. Note the 60-power telescopes on the railing of the outdoor observing platform. The entire first floor of the building is a classroom.




Fig. 1. The location of Delta Cephei.

THE VARIABLE STAR Delta Cephei is well suited to give elementary astronomy students the satisfying experience of observing a celestial object whose brightness varies from night to night.* The star is easy to find, and is sufficiently bright even at minimum; it has a moderate range in brightness and an interesting, asymmetrical light curve. A few weeks of observations usually suffice to suggest to the observer that the star goes through a regular cycle of light variation. A few months permit a determination of the period of variation and give further insight into the light curve.

The prototype of the Cepheid variables is in an extremely favorable position for observers in the Northern Hemisphere. Observations may be made in the evening during the first half of the usual academic year. By the end of January, the star gradually disappears into the evening twilight, and both the diurnal

and yearly motion of the earth cannot have failed to impress the regular observer.

Delta Cephei is easily located by means of a small isosceles triangle between Deneb (Alpha Cygni) and Cassiopeia (Fig. 1). The base of the triangle is formed by Zeta (ζ) Cephei and Epsilon (ϵ) Cephei; the vertex, Delta (δ) Cephei, points toward Cassiopeia. Zeta and Epsilon make ideal comparison stars, representing the approximate limits within which the brightness of Delta Cephei varies. The observer may adopt magnitudes for Zeta and Epsilon (for example, 3.60 and 4.36, respectively, from the Potsdam *Durchmusterung*), and from night to night assign an estimated magnitude to the variable star. A better procedure is to adopt a scale of, say, 10 steps between Zeta and Epsilon and to estimate Delta on this scale.

At Swarthmore, for several years now, we have used the following scheme. The magnitude grade A is assigned to Zeta Cephei, the magnitude grade E to Epsilon Cephei. The brightness of Delta is es-

timated on the scale A, B, C, D, E, and intermediate grades are also assigned; one-half grade is close to 1/10 magnitude. Using this grade system, there has never been any confusion as to which symbol stands for the bright and which for the faint comparison star. The variable is never estimated brighter than A; however, occasional estimates below E are made. There is some personal equation between the range in observed grades for different individuals; color equation probably plays a role. Zeta Cephei, of spectral type K0, is redder than Epsilon, of type F0, while Delta is redder at minimum than at maximum.

Starting early in October, by January an industrious student will usually have made up to 60 observations, or even more. Each evening only one observation is made, and its comparative certainty is noted in relation to the observing conditions (clear, haze, moon, and so forth). The time is recorded, perhaps to within five minutes. This is done as a matter of observing policy, rather than necessity. The times of the night may be neglected, and only the days considered, since it is quite evident that Delta does not vary noticeably during the course of one evening. The observations are plotted, using brightness as ordinate and date as abscissa (Fig. 2). Unless the number of observations is woefully inadequate, their cyclic nature should be evident after several weeks, with a number of apparent maxima and minima. As a rule, observations marked A, A-B, and possibly B, are considered as having been made near maxima. Similarly, those marked E, D-E, and perhaps D, are considered to be near minima. To analyze the observations for periodicity, a count is made of the number of days between successive maxima, and also between successive minima. These intervals cluster around values of the period and multiples thereof. An approximate value of the period may now be estimated, and it becomes possible to assign order numbers to the maxima and also to the minima, since the observer can allow for the missing maxima and minima that were not covered by the observations.

A continuous curve is then drawn, passing close to the observations, also tentatively through those parts of the plot where there are no observations but where the approximate trend of the brightness may be inferred from the cycle indicated by the observations. It should be kept in mind, however, that for a variety of causes individual brightness




Fig. 2. The preliminary light curve is dashed where observations are absent.

*See also D. B. McLaughlin, *Astronomical Journal*, 43, 113, 1934; J. Texerau, *l'Astronomie*, 65, 481, 1951.

estimates may be off, occasionally as much as one grade or more.

As the period of light variation is assumed to be constant, it is now possible to determine this length of time more exactly. The dates of the observed maxima and minima are listed, by using the days numbered consecutively from the time of the first observation, disregarding months. These numbered dates are referred to as epochs. The period may be determined by comparing the order numbers of the observed maxima and minima with the corresponding epochs. The numbers of these observed maxima are listed and paired off, using each observation only once. On the basis that the greatest accuracy is obtained by pairing two extremes in time, begin by comparing the first and last maxima to get one determination for the period; next, compare the second and the next-to-last maxima, thus obtaining another determination, and so on. The minima are handled similarly.

The conditional equations are of the form $Pn = I$, where P is the period, n the number of cycles, and I the elapsed interval. Obviously, those determinations that are based on the greatest time intervals are most accurate. In accordance with the theory of errors, we weight the observations in proportion to the square of the number of periods; putting it more technically, we derive normal equations both for maxima and for minima, as follows:

$$P \sum n^2 = \sum I n$$

The actual method of using these equations is illustrated on page 212.

Proceeding in this way, we find that values of the period derived independently from the maxima and minima usually agree within 0.1 day. If we wish, we may add the normal equations covering both maxima and minima, and thus get one simple normal equation, yielding one final determination of the period, which usually is calculated to 0.01 day, although it may be off by several hundredths of a day.


The adopted period is now used to reduce all observations to one cycle cover-

ing one whole period. Assuming that the pattern of light variation of Delta Cephei is alike in all its cycles, we may superimpose these cycles into one cycle to show their average behavior. This is done by selecting, for example, the first cycle, and reducing all subsequent observations to this first cycle. For any observation in the second cycle subtract the period from the observed epoch, and plot the observed brightnesses for the epochs thus reduced together with the first-cycle observations. For the third cycle, subtract twice the period from the observed epochs, and so on for additional

wide a time interval. The grades A, B, C, D, and E, may be replaced by the numbers 4, 3, 2, 1, 0, and averages formed of the grades (ordinates) and phases (abscissae) of all the points within each segment. The average grade and phase in each segment determines the average point, the so-called normal point for that segment. These normal points represent a condensed summary of the observations and their analysis for periodicity.

A smooth curve may be drawn which closely follows the normal points. Care should be taken that approximately equal

Fig. 3. The smoothed light curve, obtained after the observations have been reduced to a selected cycle, and after a number of normal points have been determined by averaging observations within selected intervals.



cycles. Integral multiples of the period are subtracted from the epochs of all succeeding observations, in order to place them within the first cycle, always carrying out the calculation to 0.01 day. When all observations are plotted this way, the numerous individual observations, each reduced to its corresponding "phase" in the first cycle, show the general pattern of variability (Fig. 3).

The observations normally show a healthy scatter, and there probably will be a few high residuals. The pattern of the light curve can be seen more clearly if the cycle is divided into several phase intervals, containing approximately equal numbers of observations (say anywhere from four to 10), and an average point is found within each segment. The range in phase should be sufficiently small lest the shape of the light curve be distorted by averaging observations over too

numbers of observations are above and below the final smooth curve. This procedure will confirm what may already have been indicated by observations within individual cycles, namely, that there is a difference in the slopes of the ascending and descending portions of the curve. It will also be found that the maximum brightness of the light curve is not as bright as A and the minimum not as faint as E; in other words, observations denoted A and E represent exaggerations of the true brightness range. This is not surprising, since observing errors of a whole grade or even larger do occasionally occur. The more advanced student may wish to study the distribution of the individual residuals from the "final" light curve. If desired, the period may be further improved from an analysis of the intermediate estimates, since now it can be decided which fall on the ascending and which fall on the descending branch of the light curve.

This project is an interesting experiment in observation and analysis. The student, in common with the professional, is often disappointed at the size of the residuals, and may be tempted to discard observations. With very few exceptions, which should be carefully documented, and apart from having no ethical foundation, this procedure is futile. Generally speaking, no improvement is reached by discarding large residuals; rejection of the largest residual will leave a next largest residual, and so on. As a rule, the only way to improve a series of observations of this kind is to add more observations.

(Example on page 212)




Fig. 4. A photovisual light curve of Delta Cephei, by A. J. Wesselink, reproduced from the "Bulletin" of the Astronomical Institutes of the Netherlands, 1946.



The appearance of some very bright meteors was once thought to be purely an atmospheric phenomenon. This is a 19th-century drawing of a fireball.

VISITORS from outer space? Incredible! "Physically impossible," was the conclusion of Professor P. Bertholon, one-time editor of the *Journal des Sciences*. And the accounts themselves? "Evidently untrue."

Quite so, agreed the editor of the *Décade Philosophique*: These reports are difficult to believe; one would do much better to deny their truth altogether. "Of course," you say today. "These flying saucer stories. . ."

But Professor Bertholon and his colleagues weren't talking about flying saucers at all. In fact, in the 1790's when these gentlemen made their pronounce-



E. F. F. Chladni, 1756-1827, a German physicist who revived the theory that meteorites were of extraterrestrial origin.

Stones from the Sky?

BY HELMUT HECKSCHER

ments, nothing could have been further removed from their minds than the idea of a ship from outer space. They were talking about other reports, seemingly just as ludicrous, of stones that were said to have fallen from the sky. For example, such an apparently levelheaded man as the mayor of Agen had gone so far as to affix his signature to such an eyewitness account. "How sad it is," wrote Bertholon, "to see a whole municipality verify — by means of a protocol — folklore which one can only pity. . ."

What made it worse, thought Bertholon, was that some other natural philosophers actually were beginning to take these reports seriously. In 1793, for instance, G. C. Lichtenberg, a professor at the University of Goettingen, had suggested to one of his students that certain meteoric phenomena, such as shooting stars, fireballs, and even the stones that were supposed to have fallen, might be of extraterrestrial, "cosmic" origin, and that, in any case, the topic seemed to be an interesting one for speculation. This student, a certain E. F. F. Chladni, with German thoroughness plowed through all the old and musty volumes of the *Philosophical Transactions* and other magazines that contained observations on "fallen masses."

It was all right, perhaps, for this young man to look through old papers, if he liked that sort of thing, but it was a different matter for him to publish his findings the following year. He proposed, with apparent seriousness, that masses of iron and stone had actually fallen from the sky from time to time, that shooting stars and fireballs were basically identical, and that all these bodies had arrived on earth from outer space!

Bertholon and his colleagues could only shake their heads. In ancient times, of course, there had been people like Anaxagoras who claimed that fireballs and the like were cosmic; one could regard these sayings as colorful expressions of primitive myths. But when it came to modern times, to the end of the 18th century, to the age of reason — one could only laugh at Chladni's conclusions.

But more reports began to come in. Only about two months after the publication of Chladni's findings, for instance, a great many stones were observed to fall near Siena in Italy, only about 18 hours after Mt. Vesuvius had erupted. What could be made of this, the Earl of Bristol, who had resided in Siena, asked himself?

"Either these stones have been generated in this igneous mass of clouds which produced such unusual thunder," he

wrote to Sir William Hamilton, "or, which is equally incredible, they were thrown from Vesuvius at a distance of 250 miles. . . The philosophers here incline to the first solution. . . I am reduced to perfect skepticism."

The event aroused the interest of the scientific world. To investigate the matter, Sir William himself climbed Vesuvius trying to find stones that resembled those that had fallen near Siena, but on the volcano he could find nothing but a thick coat of ashes. The astronomer, H. W. M. Olbers, too, at first thought that the stones had been hurled from the volcano, but discarded this theory when he found that none had fallen near Vesuvius, and that furthermore the Siena stones strongly resembled others that were said to have fallen in earlier times at different places.

Perhaps, Olbers speculated, stones are sometimes thrown from active volcanoes on the moon. Laplace too thought this might be possible, and calculated the

found in Siberia.

acquainted; that no power with which we are acquainted is able to give to such bodies so rapid a projectile force in a direction almost parallel to the horizon; that the matter does not rise upwards from the earth, but exists previously in the celestial regions, and must have been conveyed thence to our earth. In the opinion of Dr. Chladni, the following is the only theory of this phenomenon that agrees with all the accounts hitherto given; which is not contrary to nature in any other respect; and which besides seems to be confirmed by various masses found on the spot where they fell.

As earthy, metallic and other particles form the principal component parts of our planets, among which iron is the prevailing part, other planetary bodies may therefore consist of similar, or perhaps the same component parts, though combined and modified in a very different manner. There may also be dense matters accumulated in smaller masses without being in immediate connexion with the larger planetary bodies, dispersed throughout infinite space, and which, being impelled either by some projecting power or attraction, continue to move until they approach the earth or some other body; when being overcome by their attractive force, they immediately fall down. By their exceedingly great velocity, still increased by the attraction of the earth and the violent friction in the atmosphere, a strong electricity and heat must necessarily be excited, by which means they are reduced to a flaming and melted condition, and great quantities of vapour and different kinds of gases are thus disengaged, which defend the liquid mass to a monstrous size, till, by a still farther expansion of these elastic fluids, they melt at length burst. Mr. Chladni thinks also, that the greater part of the shooting-stars as they are called, are nothing else than fire-balls, which differ from the latter only in this, that their peculiarly great velocity carries them past the earth at a greater distance, so that they are not so strongly attracted by it as to fall down, and therefore in their

B 2

page

A facsimile reproduction of a page from the "Philosophical Magazine" of 1798, in which Chladni speculates on the origin of meteors from outside the earth's atmosphere.

LXXI. *Mémoire on the Stones said to have fallen from the Heavens. Read in the French National Institute by C. VAUQUELIN*.*

WHILE all Europe resounded with the report of stones fallen from the heavens, and while philosophers, divided in opinion on this subject, were forming hypotheses to explain the origin of them, each according to his own manner, Mr. Edward Howard, an able English chemist, was pursuing in silence the only route which could lead to a solution of the problem. He collected specimens of stones which had fallen at different times, procured as much information as possible respecting them, compared the physical or exterior characters of these bodies; and even did more, in subjecting them to chemical analysis by means as ingenious as exact.

It results from his researches, that the stones which fell in England, in Italy, in Germany, in the East Indies, and in other places, have all such a perfect resemblance that it is almost impossible to distinguish them from each other; and what renders the similitude more perfect and more striking is, that they are composed of the same principles and nearly in the same proportions.

Before the last results of the labour of Mr. Howard were known in France, I had employed myself on the same object; and I have the satisfaction to find in his memoir, which has been since printed, that they perfectly agree with those which I had obtained.

I should have abstained from any public notice of an object which has been treated of in able a manner by the English chemist, if he himself had not induced me to do so during his residence at Paris; had not the stones which I analysed been from another country; and had not the interest excited by the subject rendered this repetition excusable.

It is therefore to gratify Mr. Howard, to give, if possible, more weight to his experiments, and to enable philosophers to place full confidence in them, rather than to offer any thing new, that I publish this memoir. One of the stones which I examined was transmitted to me by C. Saint-Amans: it fell at Créon, in the parish of Juliac, on the 24th of July 1790, about nine in the evening. This stone appeared in the air under the form of a fire-ball, which was visible in almost the whole of the south of France. A very

* From the *Journal des Mines*, No. 96.

correct.

A facsimile reproduction of a page from the "Philosophical Magazine" of 1803, in which C. Vauquelin discusses the work of Edward Howard in determining the chemical composition of meteorites.

necessary initial velocity of escape.

There were more theories and counter-theories, heated discussions in the scientific periodicals. By the turn of the century a man might still deny the truth of "fallen masses," but he could no longer laugh at the reports—he must give an explanation as to how these reports came about.

Such a man was C. Patrin, who did not believe that stones had ever really fallen from the sky. He read the reports and found them very unreliable. Had there ever been a stone fall witnessed by a reputable natural philosopher? No, always farmers, natives, ignorant village people. But one could argue with Mr. Patrin—these stone falls are so rare that it would be a great coincidence if a natural philosopher happened to be present. And all the accounts agree with one another so much: Almost always a fireball is first seen to move across the sky with great speed; then there is an explosion, and finally the stones are found. A cloud is frequently seen in the sky. Furthermore, Edward Howard in England has analyzed many of these stones and has found that they resemble each other to a remarkable degree.

Mr. Patrin conceded that the observations were not incorrect, but the inferences were, and developed a theory in explanation. The rapidly moving

fireball is in reality nothing but a stroke of lightning, as evidenced by the loud report and by the cloud found to hang in the sky. When the lightning strikes the ground it melts it into a stone which is then found, still hot. The ignorant believe that something solid has actually fallen. For some reason or other lightning has a special affinity for certain earths, which it strikes in preference to others, and the likeness of the resulting stones is a necessary consequence.

This was a beautiful theory, but it had to be given up. One could point out that stones also fell on clear, cloudless days when lightning was out of the question, that no stones had ever been found where lightning was actually observed to strike. And later analyses by Mr. Howard showed that many of these stones contained nickel, an element very rare on the surface of the earth.

C. Vauquelin concluded: "The opinion which makes them come from the moon, however extraordinary it may appear, is, perhaps, the least improbable; and if it be true, that no direct proofs can be given of this opinion, it is equally certain that no well-founded reasoning can be opposed to it. The most prudent course to be pursued in this state of things is freely to acknowledge, that we are entirely unacquainted with the origin of these stones, and of the causes which produced them."

Even Patrin had to admit finally that stones had actually fallen. "The new proofs," he wrote, "leave nothing to be desired." And in the same year, 1803, the French Royal Academy was brought around to the same view by a report by

Because volcanoes are known to throw "bombs" and rocks to great heights, the fall of meteorites was often attributed to volcanoes. This volcano erupted on Lepari Island in 1888.



H. W. M. Olbers, 1758-1840, a famous German astronomer who joined in discussions of the problem of the origin of meteorites.

one of its younger members, Jean Baptiste Biot, who had been sent out to investigate a report of fallen stones.

From admitting the fall of meteorites to agreeing with Chladni as to their cosmic origin was, however, a very long step. Twenty years later the question was still undecided. Wasn't it more likely that meteorites form in our own atmosphere, in the manner of rain and hail, asked P. Egen, a German teacher



Many of the "fallen masses" that were being discussed and analyzed were first recovered by nonscientific people, in many parts of the world. This is the Guffey meteorite, found in the early 20th century by two cowboys in Park County, Colo. The composition of iron meteorites like this one, which is 88.7 per cent iron, 10.6 per cent nickel, and 0.54 per cent cobalt, was beginning to be known, but was not easily explained, in the late 1700's and early 1800's. The Guffey meteorite is three feet long and weighs 682 pounds. American Museum of Natural History photograph.

of mathematics and physics. What better proof of their earthly origin could there be than that virtually all of their constituent elements are found right here on earth?

But Chladni, by then a professor famous for his studies on sound, replied that the only conclusion was that nature in creating the universe had made use of fairly similar materials. Further, how could the atmosphere possibly carry the elements found in a meteorite?

But Egen had a theory, too: Consider the enormous quantities of solid matter that are daily volatilized in the processing of ore alone. These vapors rise to the higher regions of the atmosphere, where the electric fluid makes them

combine to form solid compounds. In addition, the electric fluid gives them enough energy to account for their high velocity and for their sometimes nearly horizontal paths with respect to the earth. The electric fluid also ignites the sulphur which they contain and thus makes them luminous while they descend.

This "electric fluid" had at the time just been propounded as a force of nature to which various electrical phenomena could be attributed. It was made to account for all and sundry. But Egen got much support, and it was found that even the famous Edmund Halley had stated in 1719 that he then believed meteorites to be of terrestrial origin.

Someone else tried to tie in meteorites with another "electric" phenomenon, the northern lights, and "proved" by means of statistics that there was a marked correlation between them.

The reader, secure in his present-day knowledge, may smile at the naivete of some of these early theories, and at the apparent blindness of the scientists of that time. He may know how in the 1830's the radiant origin of some meteor showers became recognized and how this gradually led to the acceptance of their extraterrestrial origin. But let him remember that, if our forefathers were blind, so surely are we, as we tap with ceaseless curiosity along the slowly retreating wall of the great unknown.

Student Observations of Delta Cephei — An Example

(Complete discussion by Peter van de Kamp on page 208)

AN ILLUSTRATION of the procedure is given from a series of observations made by Phoebe Burnett, Swarthmore '53, on 61 evenings from 1950 October 1 to 1951 January 11 (October 103). The observations range from A-B

to D-E, and are plotted in Fig. 2, showing the first step in the reduction procedure. From this preliminary curve the order numbers and epochs of the apparent maxima and minima are computed, as listed at the left.

The second step is to take the differences for the most extreme epochs, as for the maxima: $19 - 1 = 18$, $100 - 4 = 96$; then for the next most extreme epochs, $18 - 2 = 16$, $95 - 10 = 85$; proceeding in this manner to pair off epochs of successively shorter intervals and to set up the following equations of condition ($nP = I$):

Maxima	Minima
18 P = 96	19 P = 101
16 P = 85	17 P = 91
14 P = 74	14 P = 76
12 P = 64	11 P = 59
10 P = 53	7 P = 38
8 P = 43	4 P = 21
5 P = 26	
2 P = 11	

The third step is to apply the normalizing or weighting factors by multiplying both sides of each equation by the respec-

tive value of n . The first maximum equation thus becomes $18 \times 18 P = 96 \times 18$, or $324 P = 1728$; the second maximum equation becomes $256 P = 1360$; and so on. These equations are then added, which produces the two normal equations:

$$\begin{aligned} \text{Maxima: } 1113 P &= 5918; P = 5.32 \text{ days} \\ \text{Minima: } 1032 P &= 5529; P = 5.36 \text{ days} \end{aligned}$$

The average period of 5.34 days was used to plot the points in Fig. 3, and these were then averaged to obtain the seven normal points through which the final curve was drawn as closely as possible. The adopted period is not far from the well-established value of 5.3663 days.

The maximum brightness is somewhat brighter than B, the minimum somewhat fainter than D, representing a range of nearly half a magnitude. For comparison, Wesselink's photovisual light curve is given as Fig. 4. This curve is based on 2,448 exposures on 236 plates, taken on 159 nights in the interval 1932-1942. Wesselink's range in brightness is 0.85 magnitude; the smallness of Miss Burnett's magnitude range is probably due to color equation.

Maxima		Minima	
No.	Date	No.	Date
1	Oct. 4	1	Oct. 2
2	10	2	7
3	15	3	12
4	20	4	18
5	25	6	28
6	30	8	39
7	36	9	45
8	41	12	60
9	46	13	66
10	52	15	77
12	62	17	88
14	73	19	98
15	78	20	103
16	84		
17	89		
18	95		
19	100		

Amateur Astronomers

LEAGUE CONVENTION FIELD TRIPS AND PROGRAM

THREE FIELD TRIPS have been planned for the coming Astronomical League convention to be held in Washington, D. C., over the Labor Day weekend, September 4-7.

On Friday evening, September 4th, there will be a tour through the U. S. Naval Observatory. Located on the observatory grounds are a 26-inch refractor and 40-inch reflector, as well as other instruments. If weather permits, there will be an opportunity to view the sky with one of the large telescopes. The delegates will also be taken for a tour of the buildings, including that housing the National Capital Astronomers' 5-inch Alvan Clark refractor.

The Naval Observatory is one of the few institutions in the world and the only one in the United States at which fundamental positions of the sun, moon, planets, and stars are continually determined. From some of the observations, time determinations are made, and upon others are based the tables and star catalogues used in preparing the annual navigational almanacs. Time signals sent by radio from various government stations are regulated by the Naval Observatory.

Georgetown University Observatory will be the site of the field trip on Saturday evening, where conventionites will be guests of the Rev. Francis J. Heyden, S.J., director of the observatory. After a buffet supper served on the grounds, there will be a trip through the observatory, one of the oldest in the United States, and observing, weather permitting. Amateurs will also have an opportunity to view the photographs taken at the observatory and on the various Georgetown eclipse expeditions.

On Sunday morning, September 6th, the convention will have a field trip to Anacostia, D.C., to see the radio astronomy equipment of the Naval Research Laboratory, which consists of several radio telescopes, the largest a solid metal parabolic reflector, 50 feet in diameter. This "dish" operates on a modified gun mount atop one of the buildings. Since a gun mount is essentially altazimuth, it is necessary by automatic controls to convert the motion to that of an equatorial mounting, so as to drive the big reflector properly. The convention committee must submit a list of all persons who are to be admitted to the Naval Research Laboratory, and it will therefore be necessary to register in advance in order to make this trip.

Chartered buses will provide transportation on the field trips. The round-trip fare for the Naval Observatory and Naval Research Laboratory trips is 75 cents each, and the trip to Georgetown

Observatory is \$2.25, including buffet supper and bus fare.

Persons who are interested in attending the convention and taking part in the field trips should send the registration fee (\$1.00 until June 1st, after that \$1.50) to Mrs. Ione Alston, 20 Platts-

9:00 a.m. to noon
9:00 a.m. to 6:00 p.m.
3:00 to 6:00 p.m.
6:00 to 7:30 p.m.
7:30 to 10:00 p.m.

9:00 to 10:00 a.m.
9:00 to 10:00 a.m.
10:00 a.m. to noon
1:00 to 5:00 p.m.

6:00 to 10:00 p.m.

9:00 a.m. to noon
1:00 to 3:00 p.m.
3:00 to 4:00 p.m.
4:00 to 5:30 p.m.
7:00 to 10:00 p.m.

9:00 to 11:30 a.m.

11:30 a.m. to 12:30 p.m.

12:30 p.m.

Friday, September 4, 1953

Council meeting.
Setting up exhibits.
Registration.
Past presidents' and officers' dinner.
Field trip to U. S. Naval Observatory.

Saturday, September 5, 1953

Registration.
Middle East regional meeting.
Business session. Reports. Announcements.
Observing session: lunar, variable star, planet, and aurora sections.
Field trip to Georgetown Observatory. Buffet supper on lawn. Observing.

Sunday, September 6, 1953

Field trip to Naval Research Laboratory.
Business session. Elections.
Junior session.
Radio astronomy session.
High-altitude rockets and related subjects.

Monday, September 7, 1953 (Labor Day)

Instruments session. The latest in amateur and professional equipment.
"Operation Eclipse," planning the 1954 total eclipse expedition and convention.
Adjournment.

NORTH CENTRAL CONVENTION

The North Central region of the Astronomical League will hold its annual convention in the Quad-city area, on Friday and Saturday, July 17-18. The Popular Astronomy Club of Moline, Ill., is the host society. Joseph A. Anderer is the regional chairman.

Registration opens at 2 o'clock Friday afternoon, followed by a session in the library building of Augustana College, in Rock Island, devoted to papers and society reports. A picnic supper will be held at Carl H. Gamble's Sky Ridge Observatory in Moline, followed by observing with the 5½-inch Zeiss refractor and portable instruments. There will be a lecture Saturday morning, and a business meeting, with convention adjournment shortly after noontime.

Augustana College is providing dormitory sleeping quarters, including breakfast, at \$2.00 per person. Overnight reservations and pre-convention registration (\$1.00) may be sent to H. John Schieck, 3303 Coaltown Rd., Moline, Ill.

CHICAGO JUNIOR GROUP

The Burnham Astronomical Society of Chicago now has a junior section, consisting at present of 23 members, aged nine through 15. The group has existed independently for over a year and a half, under the guidance of Albert V. Shatzel, assistant director of the Adler Planetarium. Biweekly meetings are held, at which Mr. Shatzel lectures, and it is planned

burg Court, N. W., Washington 16, D. C., who will then forward more detailed information about the convention, hotel reservations, and the field trips.

Under the chairmanship of Charles H. LeRoy, the program committee has drawn up the accompanying tentative convention program. Sessions for papers and the exhibit will be at the Carnegie Institution of Washington.

G. R. WRIGHT, convention chairman

for members of the adult organization to give talks on their specialties, thereby providing the juniors a well-rounded program.

The telescope making facilities of the planetarium have been used for mirrors from 4½ to 10 inches, many now mounted in completed instruments. One member is making variable star observations, and efforts will be made to widen the interest in observing among the juniors.

JOSEPH A. ANDERER
7929 S. Loomis Blvd.
Chicago 20, Ill.

THIS MONTH'S MEETINGS

Dallas, Tex.: Texas Astronomical Society. June 15, 8 p.m., field meet, Dr. Arch J. McNeill's "Log Cabin," 7814 Forney Rd., public invited.


Geneva, Ill.: Fox Valley Astronomical Society. June 14, annual picnic, Aurora College, Aurora, Ill. 4:30 p.m., Wagner Schlesinger, Adler Planetarium, "Eclipsing Binaries"; 6:30, picnic supper; 8:30, observations.

Indianapolis, Ind.: Indiana Astronomical Society, 2:15 p.m., Link Observatory. June 7, Dr. Goethe Link, "The Link Observatory."

Kalamazoo, Mich.: Kalamazoo Amateur Astronomical Association. June 13, Wolf Lake. Edgar Pashby, "Lenses."

Washington, D. C.: National Capital Astronomers, 8:15 p.m., Commerce Building auditorium. June 6, astronomical films on "Viewing the Universe."

TOTAL ECLIPSE OF JUNE 30, 1954



This map, reproduced from the "American Ephemeris and Nautical Almanac," shows the vast region from which the partial phases of the total eclipse of next June will be seen, and the path of the total phase. Universal time is used.

Next Year's Favorable Total Eclipse

BY PAUL W. STEVENS

PREPARATIONS are now being made for observations of the next total eclipse of the sun. It will take place on June 30, 1954, and is of great interest to Americans and Canadians because the path of the total phase begins in the northern part of the central United States and continues in a northeasterly direction across Lake Superior and much of Canada. Readers were introduced to this eclipse in a note last August.

The eclipse should be widely observed, inasmuch as it passes over more land than water. Although much of the area is sparsely inhabited, the path of totality crosses near large centers of population, particularly in the United States, Scandinavia, and the Soviet Union. European and Russian astronomers will also undoubtedly make elaborate plans for an eclipse that may be observed with a minimum of travel time.

The circumstances of this eclipse are in many ways similar to those of the one on July 9, 1945. It is suggested that readers refer to the numerous articles in *Sky and Telescope* of that year, including

both those which gave advance information and those which recounted actual eclipse experiences. There was something from January through October.

The 1945 eclipse began at sunrise in Idaho, the shadow crossing the Montana-Saskatchewan border near Opheim, Mont., thence proceeding directly across Lake Winnipeg and out over Hudson Bay midway between Fort Severn and York Factory. Next year's shadow will begin about 800 miles farther east, in Holt County, Neb., pass across Minneapolis and St. Paul, out over Lake Superior and Rupert House at the foot of James Bay. This region is closer to large centers of population and even more accessible than was the same portion of the 1945 eclipse path.

Because it occurs on a Wednesday morning shortly after the regular school year is over, the eclipse will be very conveniently timed for most people in the United States who wish to travel to the path of totality and return during a one-week vacation. For travelers by automobile, the Keweenaw Peninsula in

Michigan, on the south shore of Lake Superior, is recommended. There the entire eclipse will take place after sunrise, and the duration of totality on the central line will be 82 seconds, with the sun and moon at an altitude of 10 degrees. The maximum duration of totality in 1945 was only 75 seconds, in Greenland; thus, this eclipse is far more favorable to observers in North America. In 1945, the sun was at an altitude of nearly nine degrees at Opheim, Mont.; in this respect, the eclipses are similar.

A total eclipse of the sun can be most favorably observed near the meridian at which mid-eclipse occurs at noon, in this case the Greenwich meridian. It is in that vicinity that the sun has its highest altitude and the duration of totality is a maximum (155 seconds, at a point west of southern Norway). Except for modifications due to climate and topography, such conditions are best for scientific observations.

When expeditions travel great distances to observe an eclipse, it is desirable to choose a site within the middle third

of the length of the central line. In this case the middle third lies mostly over water, although crossing Greenland and other important islands of the North Atlantic. Nevertheless, most of the observing will probably be done in the remaining two thirds of the shadow path, both in North America and in Europe and Asia. This will be particularly true for the geodetic measurements, as it is advantageous to station parties at points well distributed along the entire path, so measurements can be taken to tie together the fundamental triangulation surveys of widely separated geographical entities. The 1954 eclipse is particularly favorable to work of this kind and will tie together parts of three continents.

In the United States and Canada, it is hoped that the geographical distribution of amateur observers will be widespread. Of particular value in establishing the exact limits of totality are observations by those who must remain at home and will be located near the edges of the path.

As a sunrise phenomenon, the eclipse will be of unusual interest in our northern Middle West, for a total eclipse near sunrise or sunset merits considerable attention as a spectacle of grandeur. The rosy color of the sky is of great beauty when the eclipse is seen near the horizon, the varieties being as endless as those of a normal sunset. The rising of a thin

crescent sun in a darkening sky holds a thrill for the observer.

Little has been written of the view seen where totality occurs below the horizon, and it is hoped that in 1954 many such observations will be made. The changing color and brightness of the dawn follow an unexpected pattern, and the sweep of the moon's shadow across the sky is a novelty. The zodiacal light appears of unusual brightness and extent and should be of special interest where moon and sun are five to 10 degrees below the horizon. There will be fine opportunities to see this phase of the phenomenon in the Rocky Mountains, where conditions of topographic altitude and atmospheric clarity are favorable.


One of the interesting sidelights of a total solar eclipse is the appearance of stars and planets during totality. In June, 1954, the stellar background will include the famous bright stars of northern winters. A most rare circumstance will be the invisibility of Jupiter behind the solar disk. The giant planet will be obscured by the sun throughout the entire period of the eclipse for an observer anywhere in the world. This situation was discussed by the writer in *Sky and Telescope* for May, 1947.

Mercury will be a slender crescent (telescopically) close to the sun, rising for observers near the American shore of Lake Superior. Venus will make its ap-

pearance during totality in eastern Canada, while Saturn will not rise until the moon's shadow is somewhere in Europe. Mars will be close to opposition to the sun, and will be observed to disappear in the west after dawn has broken.

The writer has studied the eclipse supplement to the 1954 *American Ephemeris* describing this eclipse and mentioned in the note of August, 1952. Using data in the supplement and in the *Ephemeris* itself, computations have been made revealing circumstances of the eclipse that are of general interest. This is in addition to the information that can be found directly by reference to the tables and maps printed in one or the other of these two sources. In subsequent issues of this magazine, it is planned to reproduce some of the special maps in the supplement and to explain their important features.

NOTE: *Sky and Telescope* will act as a clearinghouse for its readers on plans and problems concerning this eclipse. Correspondence is solicited from amateur and professional astronomers and groups interested in observing the eclipse, who may wish to announce their observing plans or request information from our readers. It will be appreciated if the directors of observatories everywhere will communicate their preliminary plans for observing the eclipse as soon as these become definite enough to merit publication. Every effort will be made to give prompt publication to pertinent eclipse articles, notes, announcements, requests, and eclipse information of all kinds. —ED.



MOON'S SHADOW ON THE EARTH IN A TOTAL ECLIPSE OF THE SUN.

ASTRONOMICAL INSTRUMENTS OF QUALITY

Optical Instruments including
standard Telescopes and all
accessories.

LABORATORY OPTICAL CO.
Plainfield, N. J.

Spitz Planetarium

No. 89

has been installed at

FREDONIA
STATE TEACHERS' COLLEGE
Fredonia, New York

SPITZ LABORATORIES, INC.

5813 Woodland Avenue
Philadelphia 43, Pa.

STAR DUST

A Photographic Star Atlas

Volume No. 1. A modern star
atlas containing photographic
charts of the north polar region
to 54° north declination

175 pages, cloth bound . . . Price \$6.50

OTHER PUBLICATIONS

The Observer's Star Atlas . . . \$1.50

Webb's Atlas of the Stars . . . \$5.50

Telescopic Observations of Mars \$2.50

Write for Price List and Descriptions

HAROLD B. WEBB

145 President St., Lynbrook, N.Y., U.S.A.

SUMMER SOLSTICE COMING SOON

Also coming soon, a new set of charts now
nearly completed. Perhaps you haven't even
bought the first two sets yet. Have
you considered the concentrated study
necessary if you get them all at once? Beat
the gun! Order the first sets now and get
them partially absorbed before the new one
arrives! The new set will be all on solar
studies. The year is almost half gone, and
you are still thinking about charts and
slides, but you haven't ordered them yet!

2 slide sets — 2 chart sets	\$24.50
2 slide sets — 1 chart set	\$21.50
1 slide set — 2 chart sets	\$15.25
1 slide set — 1 chart set	\$12.25
2 slide sets . . . \$18.50	1 slide set . . .	\$9.75
2 chart sets . . . \$6.25	1 chart set . . .	\$3.50
Single slides . . . \$1.50	Single charts . . .	\$2.20
Large charts (27" x 35") each	\$3.50
(Discounts on quantities)		

We'll exchange your card for our circular.

Projector circular if you ask for it.

ASTRONOMY CHARTED

33 Winfield St., Worcester 2, Mass., U.S.A.

BOOKS AND THE SKY

OUR NEIGHBOUR WORLDS

V. A. Firsoff. Hutchinson's Scientific and
Technical Publications, Stratford Pl., Lon-
don W.1, England, 1952. 336 pages. 25s.

PRIMARILY of interest to the arm-
chair astronaut, this entertaining book
is also of value to all who have astronomy
as an avocation. Mr. Firsoff is obviously
well equipped for his topic, and he handles
it with imagination and zest. In fact, *Our
Neighbour Worlds* might be characterized
as the most optimistic and imaginative
treatment of the solar system possible
without doing serious damage to estab-
lished observational facts.

In defending habitability of the planets,
the possibility of intelligent life on some
of them, and our prospects of travel to
them, it is surprising how far a writer
like Mr. Firsoff can go and still remain
in the domain of the possible, if not al-
ways in that of the probable. The major
portion of the book is devoted to an ac-
count of present-day knowledge of con-
ditions on the earth's fellow planets, but
this is no ordinary treatment, for the au-
thor makes generous allowances in the
interpretations of physical observations.
In introducing his long and detailed chap-
ter on Mars, he states:

"Many astronomers have taken refuge
in impartially reporting the opinions of
others. Yet, laudable as the pursuit of im-
partiality may be in itself, it is seldom
very convincing, is often cramping and oc-
casionally degenerates into a merely nega-
tive attitude of pooh-poohing every find-
ing, every suggestion, without offering
anything constructive in return."

There is no hint of negative attitude any-
where in this book! In putting each
planet's best foot forward, Mr. Firsoff oc-
casionally places a great strain on credu-
lity, as when, in discussing Mercury, he
states, ". . . the rift valleys of Mercury
offer a possible foothold to life, . . ." and
then, "For all we know Mercury could be
inhabited by an intelligent race which has
succeeded in harnessing the great potential
source of energy in the intense solar radi-
ation. . . ." He admits, "This is speculation
verging on fiction, but 'life is stranger than
fiction' and we must beware of yielding
to the present vogue of peopling the skies
with desiccated and uninhabitable editions
of Arizona."

He has no intention of so doing in ap-
praising Venus as a possible abode of life.
The observational facts are, of course,
that spectroscopic analysis reveals no
water vapor nor free oxygen above the
cloud layer of Venus, but a great deal of
carbon dioxide. These facts have led con-
servative astronomers to hold forth little
hope for intelligent life on Venus. Not so
our author! He feels that there may be co-
pious supplies of oxygen and water below
the planet's clouds, neglecting the point
that if carbon dioxide is observed at heights
above the clouds, so should be the lighter
oxygen and water. He feels that organic
processes may be available, such as those
found in the anaerobic bacteria on the earth
which extract oxygen from carbon dioxide
and can produce, in this manner, water
from hydrogen-containing substances.

And so it goes. With the skill of a
debater picking loopholes in his conserva-
tive opponents' arguments, and with very
liberal allowance for possible errors of
observation and their interpretation, he
concludes, with respect to Venus, for in-
stance, "What will explorers find on
Venus? . . . I have my own vision of a
Venusian scene. . . . A pale green sky with
a very high ceiling of ribbed yellow clouds,
through which a large golden Sun shim-
mers dimly. Below is the sea of an un-
believably brilliant blue and islands cov-
ered with exuberant, multi-colored vegeta-
tion. Tall red mountains are capped with
snow."

The two earlier sections of the book,
those describing the solar system, the
astronomer's means of attack on planetary
problems, and the consideration of the
principles and state of the art of rocketry,
do not offer the author the scope for play
of imagination that the larger section just
described does.

The first section offers a creditable re-
view of the facts of our solar system and
a mature and encompassing view of the
battery of astronomical instruments con-
cerned in man's observational attack on
the planets. It includes, however, a some-
what out-of-place account of the author's
theory of the origin of the solar system.
This utilizes a supernova explosion to
carry material to planetary distances. As
with many earlier theories of origins,
nothing is said of how the planets obtained
their very large share of angular momen-
tum, or in short, what set them in motion
around the sun. This "quantity of motion"
could not have come from the radial ejection
of material from the sun. Angular
momentum is a formidable stumbling block
for makers of theories that call for the
birth of planetary material from the body
of the sun. The author appears uninformed
of the Kuiper-von Weizsaecker approaches
to the problem.

Four centrally located chapters in *Our
Neighbour Worlds* deal with space flight
and rockets. Here our author has many
more facts to deal with—the facts of
dynamics and mechanics. Here we read
much of such things as mass ratio, free
ascent, step-rockets, ordinary and atomic
fuels, electromagnetic boosters, space sta-
tions and space fuel dumps, sun mirrors,
braking ellipses, Hohmann orbits, space
suits and space hazards. The Hohmann
orbit, incidentally, is one which takes mini-
mum energy but often maximum time—
thus, 146 days to Venus and 259 days to
Mars.

Of space hazards, meteors are not con-
sidered among the most formidable; me-
teor puncture could be expected only once
every three weeks and the resulting hole
"could be mended with an ordinary bicycle
repair outfit."

With regard to sun mirrors and space
stations, the author brings up, in passing,
the old story of the "sun-gun." This was
to be a huge sodium foil concave mirror
some 30 square miles or more in area. It
was to be "anchored" in an orbit and was
to wreak havoc on civilizations below.
What appears to have been overlooked
by enthusiastic supporters of this fanciful

weapon is that to be "anchored" the sun-gun must be 22,300 miles above the earth's surface. The mirror must, therefore, have that long a focal length, and since image size depends only upon focal length, the sun's image upon the earth's surface would be some 200 miles in diameter! With so huge an image, the light gathered by this solar mirror would be spread exceedingly thin. The mirror itself would appear to the observer below as a bright point of light less than a minute of arc across.

The critical reader will find many spots of loose thinking, of exposition simplified to the point of ambiguity, and few of downright error. Nonetheless, this book, taken as a whole, can be recommended for the armchair amateur. The illustrations are good, the paper and printing are excellent, and especially valuable is the list of references at the end of each chapter. These are indeed redeeming features. Another is a short mathematical appendix.

J. ALLEN HYNEK
Ohio State University

ADVANCES IN GEOPHYSICS, I

H. E. Landsberg, editor. Academic Press, Inc., New York, 1952. 362 pages. \$7.80.

THERE IS no real dividing line between the earth sciences and astronomy. Their close historical association is evident upon looking into any astronomical periodical from early in the last century, where it will be found that as many papers are devoted to geodesy, terrestrial magnetism, and meteorology as to astronomy itself. Today the study of the upper atmosphere is pursued equally by astronomers and geophysicists. From the astronomical viewpoint, geophysics might be defined as a special branch of astrophysics which treats the most accessible of all astronomical bodies, the earth.

Thus, there is a real need for a concise

review of current developments in the geophysical sciences, presented in a form suited to the needs of a reader whose bent is primarily astronomical. This need is well met by the new series, **Advances in Geophysics**, of which the first volume has just appeared under the editorship of H. E. Landsberg, director of the Geophysics Research Directorate, Air Force Cambridge Research Center.

Of the eight contributions in this volume, that of most astronomical interest is "Exploration of the Upper Atmosphere by Meteoritic Methods," by F. L. Whipple, of Harvard Observatory. This is an excellent 50-page summary of present knowledge about meteors, clearly enough written to be easily read by an advanced amateur. One point brought out by Whipple should be mentioned. While visual observations of meteors have been driven into obsolescence by the more powerful photographic and radar techniques, there is an important exception. This is observation of the drift of long-enduring meteor trains, from which valuable information can be gained on winds in the upper atmosphere.

The remainder of the book is more likely to attract the professional rather than the general reader. The first two chapters will particularly interest astronomers. All who are concerned with the measurement and reduction of photographs on a large scale find a serious bottleneck in the recording and processing of data. J. C. Bellamy gives a useful and thought-evoking analysis of the mechanization of analogous procedures in geophysics. Arnold Court describes new statistical techniques in geophysics which could easily be adapted to problems in astronomy.

The value of **Advances in Geophysics** is enhanced by good indices and full bibliographies. An attractive feature is the listing at the end of each chapter of the mathematical notation used.

JOSEPH ASHBROOK
Yale University Observatory

SKY PUBLICATIONS

SKY SETS—I

Pictures of solar system and galactic objects, 24 in the set, printed on heavy white paper, and suitable for study and framing. The set, \$5.50

MAPPA COELESTIS NOVA

The northern sky to -45° is shown in this large chart, and each star is colored according to its spectral class. The chart makes a fine transparency, 28 inches square. \$4.00

SKALNATE PLESO ATLAS OF THE HEAVENS

Sixteen charts cover the entire sky to magnitude 7.75: 1950 co-ordinates. Each chart area is $15\frac{1}{2}$ by $23\frac{1}{2}$ inches. \$5.50

MOON SETS


Eighteen pictures, nine at first quarter and nine for the last quarter, each on a sheet of heavy stock 12 by 18 inches. There are key charts for named lunar features. \$2.50

WORLD WIDE PLANISPHERE

A book of basic constellation charts and masks to show the stars above the horizon at any time of night or year, at all latitudes, northern and southern. By William H. Barton, Jr. \$3.00

Making Your Own Telescope
— Allyn J. Thompson \$4.00
Splendors of the Sky 50 cents
The Nature of Cosmic Rays 50 cents
Relativity 50 cents

SKY PUBLISHING CORPORATION
Harvard Observatory, Cambridge 38, Mass.



A Scientific Inquiry

By KENNETH HEUER

Illustrated by
CHESLEY BONESTELL


Can our world be totally destroyed? How? When? Can we do anything about it?

THE END OF THE WORLD explores the scientific possibilities among the ways the world actually can end:

- comet collisions
- moon, asteroid and star collisions
- the death of the sun
- the explosion of the sun
- atomic war

ALBERT EINSTEIN says: "Very good . . . rich in ideas and offers much solid knowledge in an easily digestible and very attractive form."

KENNETH HEUER is a fellow of the Royal Astronomical Society, former lecturer at the Hayden Planetarium, author of *Men of Other Planets*.



Just published
\$3.00

TEN DAYS'
FREE
EXAMINATION

RINEHART & COMPANY, Dept. ST-6
232 Madison Avenue, New York 16, N. Y.
Please send me *THE END OF THE WORLD* by Kenneth Heuer for 10 days' free examination. At the end of 10 days I agree to remit \$3.00 plus postage or return the book postpaid.

NAME
ADDRESS
CITY ZONE STATE
SAVE MONEY: Enclose \$3.00 with order, we pay postage. Same return privilege.

THE COMETS AND THEIR ORIGIN, R. A. Lytleton, 1953, Cambridge University Press. 173 pages. \$5.00.

A discussion of the properties of comets, dynamical and physical, is followed by the presentation and development of the author's theory for their origin and formation, in brief, a process "of the accretion of interstellar dust through the gravitational action of the sun during passages through galactic dust clouds." The formation of comet tails is then considered.

PHOTOGRAPHY AND THE AMATEUR ASTRONOMER, Gerald Merton, 1953, British Astronomical Association, 303 Bath Rd., Hounslow West, Middlesex, England. 24 pages. 2s.

Issued as BAA Reprint No. 1, this booklet contains the address of the retiring president first printed in the BAA *Journal* of December, 1952. (See In the Current Journals, *Sky and Telescope*, March, 1953, page 128.) The author gives a historical sketch, discusses emulsions, photogenic response, reciprocity failure, limiting magnitudes, complicating factors, diameters of star images and magnitudes, cameras. A short section on photographic work in the BAA observing sections and a full bibliography conclude the booklet, which is illustrated with halftones and diagrams.

UNITRON Leads in Value and Quality

The New 4-Inch Equatorial Heads a Distinguished Line of Nine Refractors

The choice of leading universities and active amateurs the world over! Precision made from the finest of materials by one of the world's largest manufacturers of optical instruments.



2.4-INCH EQUATORIAL

The complete instrument for the active amateur.
UNITRON Model 128

Objective: 62-mm. (2.4") diam., 900-mm. (35.5") focal length, $f/15$.
Eyepieces: 100x, 50x, 35x included.
150x: \$14.75 extra.
129x: \$8.95 extra.
72x: \$6.75 extra.

COMPLETE with equatorial mounting and slow-motion controls, tripod, view finder, star diagonal, erecting prism system, sunglass, eyepieces, wooden cabinets.

Only \$225

2.4-INCH ALTAZIMUTH

UNITRON Model 114

Not illustrated. Same optical specifications, eyepieces, and accessories as the 2.4-inch Equatorial, but with altazimuth mounting.

Only \$125

ACCESSORIES

VIEW FINDER: 42-mm. (1.6") coated achromatic objective. 10x crosshair eyepiece. Furnished with mounting brackets. A wonderful accessory for your present telescope. Only \$18.00

GUIDE TELESCOPE: 62-mm. (2.4") coated achromatic objective, 700-mm. (27.5") focal length. 78x crosshair eyepiece. Furnished with mounting brackets. Convert your present telescope for astrophotography. Only \$75.00

EQUATORIAL MOUNTING

The equatorial mounting and tripod of the 3" equatorial, Model 142, is available separately. Full specifications on request. An exceptional value.

Only \$198

We welcome your inquiries.

Illustrated literature on all instruments is available at no cost or obligation. If you have any questions regarding the instrument most suitable for your purpose, we shall be glad to give them our prompt and careful attention.

Please address your card or letter to the attention of Dept. TA-6
Microscope catalogue also available.

• LENSES are FULLY CORRECTED for spherical and chromatic aberration and are COATED for maximum brilliance and clarity of image. AIR-SPACED CELL insures freedom from "clouding" with age.

• EYEPIECES of the HIGHEST QUALITY . . . Orthoscopic, Achromatized Ramsden, Huygens.

• FINEST MATERIALS throughout. DURALUMINUM TUBE. Moving parts of BRASS carefully machined to close tolerances, and finished in CHROMIUM.

• MODERN DESIGN based on time-tested engineering principles. HANDSOME APPEARANCE to which no illustrations can do justice.

• STURDY TRIPOD may be folded for convenient storage.

• EQUATORIAL MODELS have slow-motion controls for both declination and right ascension.

• ALTAZIMUTH MODELS have slow-motion controls for both altitude and azimuth.

• VIEW FINDER with crosshair eyepiece gives wide field.

• RACK-AND-PINION FOCUSING.


• STAR DIAGONAL for easy observing at all altitudes.

• SUNGLASS for solar observation.

• ERECTING PRISM SYSTEM for TERRESTRIAL OBSERVATION may be used with any of the eyepieces to give the same complete range of terrestrial magnifications as for celestial observation.

• FITTED WOODEN CABINET, handsomely finished, for compact storage of telescope and accessories. Separate case for the tripod.

Unbelievably low prices! Outstanding features, performance, and workmanship impossible to duplicate even by paying considerably more.



3-INCH EQUATORIAL

Ideal for amateur research and school observatories.
UNITRON Model 142

Objective: 78-mm. (3") diam., 1,200-mm. (47.2") focal length, $f/16$.
Eyepieces: 200x, 133x, 96x, 67x, 48x included. 171x: \$8.95 extra.

COMPLETE with equatorial mounting and slow-motion controls, tripod, setting circles and verniers, view finder (30-mm., 6x), sun projection screen, sunglass, star diagonal, erecting prism system, eyepieces, pocket eyepiece case, wooden cabinets.

Only \$435

3-INCH PHOTOGRAPHIC EQUATORIAL

UNITRON Model 145: Not illustrated. Same specifications as Model 142 but with 78x, 62-mm. guide telescope and 10x, 42-mm. view finder.

Only \$550

3-INCH ALTAZIMUTH

UNITRON Model 140: Not illustrated. Same optical specifications and accessories as the 3-inch Equatorial but with altazimuth mounting. 171x, 133x, 96x, 67x, 48x eyepieces supplied. Sunscreen not included, but available as accessory.

Only \$265

4-INCH ALTAZIMUTH

UNITRON Model 150: Not illustrated. Same optical specifications, eyepieces, and accessories as the 4-inch Equatorial, but with altazimuth mounting. Sunscreen not included, but available as accessory.

Only \$465


EASY PAYMENT PLAN

UNITRON refractors may be purchased on an easy payment plan. Write for details.

All Instruments Fully Guaranteed
Send check or money order for full amount or 25% deposit with balance C. O. D.

Telescopes shipped Express Collect.

It's Here!



4-INCH EQUATORIAL REFRACTOR

UNITRON Model 152

Objective: 105-mm. (4.13") diam., 1,530-mm. (60.24") focal length, $f/15$.

Eyepieces: 255x, 219x, 170x, 122x, 84x, 61x. Higher and lower powered eyepieces will be made available.

COMPLETE with equatorial mounting and slow-motion controls, tripod, setting circles and verniers, view finder (42-mm., 10x), sun projection screen, sunglass, star diagonal, erecting prism system, eyepieces, pocket eyepiece case, wooden cabinets.

Only \$785

4-Inch Photographic Equatorial also available.

UNITED SCIENTIFIC CO. 204-6 MILK ST. BOSTON 9, MASS.

GLEANINGS FOR ATM's


EDITED BY EARLE B. BROWN

INEXPENSIVE PORTABLE PLANETARY TELESCOPES

THIS ARTICLE is written for the amateur astronomer who wants to build a telescope at minimum cost. The optical parts, which consist of a 3-, 4-, 5-, or 6-inch diameter mirror, ground, polished, and aluminized, ready to use, with an aluminized diagonal and two eyepiece lenses, can be purchased for a reasonable sum.

The instruments can be made at home with simple hand tools by the average home worker. The mounting is light and easily portable; it can be taken indoors when not in use, or can be carried in a car on trips to the country. Setting scales are provided for right ascension and declination, so that Venus or Mercury can be located in the daytime, when they are best seen, and comets or other objects invisible to the naked eye at night can also be found. To keep the telescope accurately trained on an object, a slow-motion manual-drive tangent screw is used. A smooth slow-motion drive is especially needed for high powers, when the object appears to slide rapidly out of the field of view. For planetary work the primary mirror should have a focal length at least 10 times its diameter. A long focal length mirror produces a larger image, so that low-power eyepieces can be used for all general work.

High-power telescopes magnify any motion or vibration, and with a flimsy mounting stars may dance around in the field, causing eye strain and making it difficult or impossible to see fine detail. To eliminate this trouble the mountings have a distance between polar-axis bear-




A view of the right-ascension tangent screw, time scale, and storage place for eyepieces, looking down the length of the carriage that forms the polar-axis support.

ings of 10" for the smallest reflector and 22" for the largest.


In declination the telescope covers a section of the heavens from the southern horizon to the zenith. At 40° north latitude the telescope can be set in declination from -50° to +40°. This section provides for all the planets, and most of the other popular objects for small telescopes.

A tube to mount the optics is required. Duro-board cores, used by paper mills, are about 3/10" thick and make very rigid tubes. These are preferable to metal tubes,



The author here demonstrates the slow motion in right ascension of the 4-inch telescope, which is one of the battery of instruments pictured on the front cover, ranging in size from 3-inch to 6-inch, all of the same fundamental design.

UNUSUAL OPTICAL BARGAINS



AMAZING Double Utility Combination 15X TELESCOPE and 35X ERECTING MICROSCOPE

Extra utility by removing telescope portion and using as 35 power microscope to inspect, dissect, etc. Telescope objective lens is achromatic, coated. Telescope can be removed

and carried in pocket — focuses from 10 feet to infinity. Microscope has 3-element, color-corrected objective — image is erect, just as your eye sees it, not upside down as in an ordinary microscope. Ramsden eyepiece with coarse or fine focusing. Gives you 3" of working distance when used as microscope. Entire unit folds down to 2" x 7" x 10". Telescope instantly adjustable in height from 3" to 14".

Stock #70,017-Y \$29.50 Postpaid

NEW HUYGENS TYPE ASTRONOMICAL EYEPIECES — Imported! Mounted! Fine quality! precision machined throughout. Standard 1 1/4" outside diameter.

Stock #30,063-Y F.L. 6 mm. (1/4") \$7.50 Patpd.

Stock #30,064-Y F.L. 12.5 mm. (1/2") 6.00 Patpd.

Stock #30,065-Y F.L. 10 and 20 mm. 8.00 Patpd. (You have a choice of 2 powers by rearranging lenses. Directions included. Chrome and black finish.)

IMPORTED MICROSCOPE

100, 200, 300 Power

Good optical qualities. Fine focusing. Certainly not the equal of a \$200 instrument — but definition is surprisingly clear and good... in fact amazingly so at this price. Serviceable construction. 1 Ocular, 3 Objective Lenses. Rack & Pinion Focusing. Revolving Disc-Light Adjustable Mirror. Square Stage (2 1/4" x 2 1/4") with Slide Clamps. The greatest microscope bargain on the market!



TRY IT FOR 10 DAYS... If you're not completely satisfied your money will be refunded in full. Instrument comes packed in sturdy, hardwood case. Stock #70,008-Y \$14.95 Postpaid

NEW 30 POWER TELESCOPE COMPLETE WITH TRIPOD

Unusual Bargain Price. A beautifully made, sturdy, imported instrument with dozens of uses. Spots 22-caliber bullet holes at 200 yds. You can see the craters on the moon! Excellent for amateur astronomers. All metal body — focuses from approx. 40 ft. to infinity. Achromatic objective — 40 mm. dia. — outside surface low-reflection coated. Smooth slide focusing eyepiece. Eye-lens outside surface coated. Lens erecting system. Images right side up. Sturdy tripod 8 1/2" high. Can be used also for camera. Maximum length 26 1/4". Barrel diameter approximately 1 1/4". Weight 1 1/2 lbs.



Stock #70,018-Y Only \$21.95 Postpaid

MOUNTED ERFLE EYEPIECE

Wide angle! Used on astronomical telescope. Consists of three achromats perfect and coated, in a metal focusing mount. Diameter of lenses 46, 46, and 38 mms. Government cost about \$54.00. Focal length 1 1/4" (32 mm.). This eyepiece will give you an unusually wide field.

Stock #5160-Y \$8.50 Postpaid

CROSSLINE RETICLE — Dia. 29 mm.

Stock #2036-Y 50c Postpaid

TELESCOPE EYEPIECE — Consists of 2 Achromatic lenses F.L. 28 mm. in a metal mount.


Stock #5140-Y \$4.50 Postpaid

Same as above but about 1 1/4" extension has been added with O.D. of 1 1/4", which is standard for astronomical telescopes.

Stock #5223-Y \$5.25 Postpaid

ASTRONOMICAL MIRROR MOUNTS

Cast aluminum with brass mounting and adjusting screws and mirror clamps. Two sizes: for 6" mirrors — mount will fit 7" I.D. tube:



Stock #50,010-Y \$7.00 Postpaid

For 4 1/4" mirror — mount will fit 5" I.D. tube.

Stock #50,011-Y \$5.25 Postpaid

We have Literally Millions of WAR SURPLUS LENSES AND PRISMS FOR SALE AT BARGAIN PRICES... also numerous low cost imported instruments! Write for Catalog "Y" — Free!

Order by Stock No. — Send Check or M.O. Satisfaction Guaranteed

EDMUND SCIENTIFIC CORP.
BARRINGTON, NEW JERSEY



WAR SURPLUS BARGAIN

OPTICS

ACHROMATIC TELESCOPE OBJECTIVES — Perfect Magnesium Fluoride Coated and cemented Gov't. Surplus lenses made of finest Crown and Flint optical glass. They are fully corrected and have tremendous resolving power and can be readily used with 1/4" F.L. eyepieces. Guaranteed well suited for Astronomical Telescopes, Spotting Scopes, etc. Gov't. cost approximately \$100.00.

Diameter	Focal Length	Each
54 mm (2 1/8")	300 mm (11.81")	\$12.50
54 mm (2 1/8")	330 mm (13")	12.50
54 mm (2 1/8")	390 mm (15.356")	9.75
54 mm (2 1/8")	508 mm (20")	12.50
54 mm (2 1/8")	600 mm (23 1/2")	12.50
78 mm (3 1/16")	381 mm (15")	21.00
81 mm (3 3/16")	622 mm (24 1/2")	22.50
83 mm (3 1/4")	660 mm (26")	28.00
83 mm (3 1/4")	711 mm (28")	28.00
83 mm (3 1/4")	876 mm (34 1/2")	28.00
83 mm (3 1/4")	1016 mm (40")	30.00

We can supply ALUMINUM TUBING for the above lenses.

SYMMETRICAL EYEPIECE LENS SET

Each set consists of two magnesium-fluoride coated and cemented achromats, and exact Gov't. spacing diagram. Gives wide flat field. 1/2" E.F.L. (20X) Lens Set 13-mm dia. \$4.50
1/2" E.F.L. (13X) Lens Set 18-mm dia. \$3.50

Rectangular Magnifying Lens — Seconds, sells for \$6.50. Size 2" x 4" \$ 1.00
First Surface Mirror 14"x16" 1/4" thick .. 10.00
First Surface Mirror 8"x10" 1/4" thick .. 4.25
First Surface Mirror 4"x4" 1/4" thick .. 1.50
First Surface Mirror 1 1/2"x1 1/2" 1/16" thick .. .25
Optical Peep Sight — Use as camera viewfinder, etc. Dia. 1 1/2", weight 1 1/2 oz. ... \$1.00

LENS CLEANING TISSUE — 500 sheets 7 1/2" x 11". Bargain priced at only \$1.00

RIGHT ANGLE PRISMS

8-mm face .. ea. \$.75 28-mm face .. ea. \$1.75
12-mm face .. ea. .75 38-mm face .. ea. 2.00
23-mm face .. ea. 1.25 47-mm face .. ea. 3.00

!!!NEW LOW PRICES!!!

BEAUTIFUL IMPORTED BINOCULARS
Precision made, at a low price within the reach of every man's pocketbook. Complete with carrying case and straps.

6 x 15 Ind. Focus .. Coated	\$16.00*
8 x 25 Center Focus ..	23.30*
8 x 30 Center Focus ..	27.50*
7 x 35 Center Focus .. Coated	41.50*
7 x 50 Ind. Focus .. Coated	35.00*
7 x 50 Center Focus .. Coated	39.25*
10 x 50 Ind. Focus .. Coated	45.00*
16 x 50 Center Focus .. Coated	55.00*

*Plus 20% Federal Excise Tax

MOUNTED EYEPIECE has two lenses 29 mm in diameter. Cell fits 1 1/4" tube. 1 1/4" E.F.L. (8X) \$4.50



Gov't. Cost \$200.

Plain Optics \$6.50 Coated Optics \$10.50

"MILLIONS" of Lenses, etc.
Free Catalogue


We pay the POSTAGE — C.O.D.'s you pay postage. Satisfaction guaranteed or money refunded if merchandise returned within 10 days

A. JAEGER'S 6915 Merrick Road
Lynbrook, N. Y.

which conduct disturbing heat waves along their inner surface, causing distortion of the image. Any material which is an electrical insulator is also an insulator of heat waves, so that cardboard, fiber, wood, or bakelite tubes can be used to advantage. If a metal tube is used, be sure to insulate the inside surface with cork or other suitable material; otherwise the telescope may have to be left outdoors a half hour or more before it settles down, and on certain nights when the temperature is continually changing, it may never completely settle down. It is the writer's experience that aluminum, which is a very good conductor, is about the worst metal to use.

For neatness and precision in mounting the optical parts, the front end of the duraluminum tube should be squared up using a fine-tooth hack saw and trimming any fuzzy edges with a knife or razor blade. Before cutting, draw a pencil line around the tube by using a piece of stiff un-wrinkled paper wrapped around the tube to act as a guide for the pencil. The paper should be at least 12" wide and long enough to go completely around the tube so that the ends can be lined up. The marking edge of the paper must be trimmed on a straight line, the exact size and shape of the paper being unimportant. Where the paper strip overlaps, a mark should be made, the paper stretched flat, and the distance around the tube accurately divided into four equal parts. One inch down from the end of the tube draw a second line and mark the tube opposite the four marks just made on the paper. This will locate the 3/16" holes for mounting the spider. On a line perpendicular to the front end of the tube and 3" below one of the holes for the spider, make a mark for the location of the eyepiece holder. A little time and patience will enable the worker to cut a neat hole 1 3/8" in diameter around this point.


The collar is made from an old piece



Compare this view of the mounting and the plumb line used for leveling with the diagram on the opposite page.

of 1/4" thick leather belting trimmed to a width of 1". This collar is used to keep the telescope tube from sliding down in the cradle. Drill holes for 3/16" by 5/8" round-head stove bolts, the first hole being 3/8" from the end of the belt and the other holes evenly spaced around the tube approximately 3" apart.


Paint the inside surface of the tube with one coat of flat black paint, and the outside with two coats of light gray enamel. All interior parts of the optical system should be painted with the flat black, being sure not to get any on the mirror or diagonal. Avoid touching the flat black as greasy fingers will spoil the dead black surface, causing reflections. (To strengthen the tube, before applying any paint give the cardboard one or more coats of varnish or shellac. This will act as sizing, and the paint will not be so readily absorbed and will spread on and dry more evenly. The more ambitious amateur will find that wooden rings turned to fit the top and bottom ends of the tube will also help strengthen the cardboard and add trim to the finished tube.)



This diagram, showing the location of the optical parts inside each tube, illustrates the manner of scaling the dimensions from the smallest up to the largest.

The telescope rests in a cradle, stopping against the leather collar, and is tied in place by a piece of lightweight sash cord. Be sure to make the cradle wide enough so that the telescope will fit freely between the side members. The bottom and sides are made of $\frac{1}{4}$ " masonite, bolted to $\frac{3}{8}$ " by $1\frac{1}{4}$ " wood strips at the inside corners with $3/16$ " stove bolts. The declination-axis pivots are two $\frac{3}{8}$ " fine-threaded hexagon-head cap screws, with a tight fit in the cradle and a free bearing fit in the carriage. The cradle carries a declination scale index graduated in degrees from 0 to 10, together with a 2" by 3" masonite clamp plate. A piece of felt under the clamp plate will prevent scratching the carriage graduations.

The carriage supporting the cradle has two side wings graduated every 10° of declination. The east and west wings carry identical graduations which can be



A diagram of the various parts of the mounting designed by Mr. Lawrence.

easily read for all positions of the telescope. The wings, bottom, and back of the carriage are all made from $\frac{1}{4}$ " masonite, the inside joints bolted together with $\frac{3}{8}$ " square wood strips using $3/16$ " stove bolts. The width and height of the back plates are 8 x $8\frac{1}{2}$, $8\frac{1}{2}$ x $9\frac{1}{2}$, 9 x $10\frac{1}{2}$, and $9\frac{1}{2}$ x $11\frac{1}{2}$ inches for the four sizes of telescopes. The bottom carries two $\frac{3}{4}$ " hardwood polar-axis bearings. The polar axis is made from $\frac{3}{8}$ " drill rod or cold rolled steel, held in place with sheet-metal straps. Note that the polar axis is stationary and the carriage bearings turn with the instrument in right ascension.

The pedestal has a central vertical plate with the polar axis mounted on its upper edge, a pear-shaped hole plate, and a front vertical plate for cross bracing, all made from $\frac{1}{4}$ " masonite. These three members are fastened together with $3/16$ " stove bolts. The hole plate is used for setting and clamping the right-ascension arm, while the front vertical plate is hinged at the bottom edge to the base plate so that the mount can be leveled. This vertical plate is also slotted for two east-west adjustable legs used for cross-leveling. A plumb line is mounted on the side of the central vertical plate with two steel brackets to indicate a true vertical

adjustment of the mount, in both vertical planes.

The tangent screw provides a smooth slow-motion drive for the telescope in right ascension, one revolution of the hand wheel being approximately one minute of time. The lead screw is made from $\frac{3}{8}$ " diameter, 16 threads per inch, standard threaded rod, which can be purchased from any large hardware store. It has a 3" masonite hand wheel on each end, with a hand crank made from any suitable radio knob. The hand wheels are clamped in place between two hexagon nuts which are also adjusted to remove any end play from the lead screw. The screw rotates on ball bearings and carries a traveling swivel nut with an index line to read against the time scale. The swivel nut is the most difficult part to make. It should be reamed for a sliding fit on the $\frac{3}{8}$ " by 6" steel rod, and carefully tapped to fit the lead screw, to work freely with a minimum of lost motion. A 2" masonite arm is held by a clamp using a $5/16$ " carriage bolt and wing nut. The arm provides a coarse setting for right ascension, and stops against a pin set in the hole plate, the holes being spaced one hour apart.

A star and planet finder, illustrated, has dials made from white-faced cardboard or bristol board with a maximum diameter of 12". The day-of-month scale is mounted to a backing of $\frac{1}{4}$ " masonite and remains fixed. The outer dial is 12" in diameter and shows the months and the hours of right ascension from 0 to 24. The second dial is $10\frac{1}{2}$ " in diameter and is graduated with a scale showing minutes of right ascension and minutes of clock time from

BERAL COATINGS

The following prices now apply on BERAL coatings for telescope mirrors:

3" diam.	\$2.50	8" diam.	\$4.50
4" diam.	2.75	9" diam.	5.50
5" diam.	3.00	10" diam.	6.50
6" diam.	3.50	11" diam.	8.50
7" diam.	4.00	12 $\frac{1}{2}$ " diam.	9.75

All prices f.o.b. Skokie, Ill.
Minimum order \$2.00

BERAL coatings are not overcoated with fluorides, quartz or silicon monoxide. They contain no chromium, and as a result they can be easily removed without danger of harming the glass supporting surface. The efficiency of BERAL coatings is uniformly higher, so that mirror performance is at its best.

LEROY M. E. CLAUSING

8038 MONTICELLO AVE. SKOKIE, ILL.

BUILD YOUR OWN TELESCOPE

KITS: Complete with instructions, high quality materials, supplied at low prices.

Plate — \$4.00 up Pyrex — \$5.50 up

Lenses Cells
Prisms Eyepieces

Mirrors ground to your order.

Aluminizing — with the new Quartz Coating

Satisfaction or Money Refunded

Write for Free Catalog


GARTH OPTICAL COMPANY

130 Garth Rd. Scarsdale, N. Y.

ORTHO SCOPIC OCULARS

WE HAVE IN STOCK for immediate delivery the best eyepiece ever produced for astronomical and scientific work, with the following characteristics:


1. Four-element design giving a flat, beautifully corrected field of 50° covering more than 160 per cent of the area of the conventional Ramsden for the same power. This eyepiece is a "must" for RFT work.
2. Precision optical elements, magnesium fluoride hard coated, increasing the light transmission approximately 10 per cent.
3. Simple take-down for cleaning.
4. Precision metal parts black anodized for anti-reflection and ground to $1\frac{1}{4}$ " O. D.
5. Clean mechanical design permitting comfortable observation and ease of focusing.



Price postpaid, \$15.95 each

These eyepieces are produced in 8 mm., 16 mm., and 32 mm. effective focal lengths only.

ASTRONOMICAL OBJECTIVES



These objectives are sold with cells and rigidly tested on double stars for resolving power before being sold.

Unconditionally guaranteed — Immediate delivery

CHESTER BRANDON

WE CAN SHIP from stock quality astronomical objectives with the following features:

1. Superior optical design utilizing a larger air separation. Color corrected on C and F and hand corrected on spacing to reduce residuals to a minimum. Completely free of coma.
2. Glass-air surfaces magnesium fluoride coated, increasing light transmission approximately 10 per cent.
3. Quality optical glass precision annealed and held to one ring test plate match.
4. Cell made to precision tolerances and suitably coated to prevent stray light reflections. Each cell engraved with effective focal length and serial number.

These objectives are supplied as follows:

3" C.A.	45" E.F.L.	\$ 49.00
4" C.A.	60" E.F.L.	\$119.00

California residents, add 5% sales tax
Box 126, Montrose, California

MAKE A PROFESSIONAL Astronomical or Terrestrial TELESCOPE for as little as \$7.00

Now you can afford the luxury of a real telescope. See the mountains of the moon, Saturn's rings, nebulae and a host of heavenly wonders. We furnish the reflecting Mirror accurately ground, polished, and aluminized, together with a flat mirror and three lenses to make two eyepieces. Other parts are usually found around the home or purchasable from us very reasonably. Thousands of satisfied amateurs have built these 'scopes. So can you! Sharp, clear images assured. No fuzziness or rain-bows.

3" Kit...\$ 7.00 5" Kit...\$17.00
4" Kit... 12.00 6" Kit... 20.50
Send for your telescope kit now. Free instructions with kit. "Know-How" Catalog, 30c.
Or buy a complete instrument from us:
Stellarscope 3" (40x & 80x)\$16.50
Deluxe Model 3" (40x & 80x) 35.00
Stellarscope 4" (60x & 120x) 26.00

EYEPIECES

Save Money — Make Your Own

It's simple. Use our All-in-One Hastings-Ross Achromatic Triplet lens system. You can hold these complete lens systems in simple bushings and have the equal of the world's finest eyepieces. Save money too.
20x or 1/2" E.F.L. Triplet Lens .. Only \$6.80
33x or 1/3" E.F.L. Triplet Lens .. Only 6.80
40x or 1/4" E.F.L. Triplet Lens .. Only 6.80
50x or 1/5" E.F.L. Triplet Lens .. Only 10.00

TREMENDOUSLY WIDE FIELD 3-Element Achromatic Eyepiece

Almost all gone. These are not rejects.

Six lenses! Finest eyepiece ever made anywhere. Our greatest buy. Made of 3 separate achromatic elements. All outside surfaces fluoride coated. In focusing mount. 1-13/16" (43 mm.) clear aperture, flat field to edges. Focal length 1 1/4" (32 mm.) (8x), 69° angle. Outside diameter of mount 2 1/4" (54 mm.). Each \$15.00 plus postage. The above with bushing to fit standard 1 1/4" eyepiece tubes. \$18.00

PORRO PRISMS

Oval Face. Manufactured by world-famous opticians for use in Army and Navy binoculars. Rejected for slightly chipped edges. Outstanding bargain!
6 x 30...50c 7 x 50...50c Giant size...75c
Add 5c per prism for packing and P.P.

POLAROID IN GLASS

Has been used successfully in Quartz Polaroid filter for observing the prominences on the sun. Experiment with polarized light. Get two genuine Polaroids that can be used in photography, petrology, physics, chemistry, astronomy, and so forth. Mounted in optical glass 3" diameter (72 mm.). In mounting, per pair Only \$2.35 P.P.

PRISM — LIGHT FLINT GLASS

Water-white glass of excellent quality. Government inspected. 45-90-45 degrees. Fluoride coated in mounting of aluminum-magnesium alloy. Especially designed for Foucault test setups. 9/16" x 1 1/16" face (15 mm. x 18 mm.). Free from scratches, abrasions, and imperfections. No chips or roughs. Use in any type optical system — photography, telescopes, etc. Weight each 1 oz. A \$15.00 value.
Only \$2.00 each

REFRACTING ASTRONOMICAL TELESCOPE

With fully corrected 60-mm. (2-3/8") diameter achromatic objective of 800-mm. focal length, which is air-spaced and coated. Made in Japan of substantial construction with all white enameled body. Focusing by rack and pinion. Very good quality and design. The eyepieces work with the erecting prism for terrestrial use, as well as with the diagonal prism for astronomical use.

SPECIFICATIONS: Altazimuth mounting; tripod of wood, with 3 collapsible sections, fitting into case, 4x finder with cross-lines; diagonal eyepiece, erecting-prism eyepiece, sunglass. Three corrected eyepieces giving powers of 40x, 64x, and 88x. Comes complete in hinged, covered wooden case. Save money! Our price only \$97.00, plus \$3.00 for packing and parcel post.

Include postage — Remit with Order

No open accounts — No C.O.D.

Send 25c for Catalog.

HARRY ROSS

TELESCOPES MICROSCOPES
Scientific and Laboratory Apparatus
70 West Broadway, Dept. ST-6B
New York 7, N. Y.

0 to 60. The third dial is 9 1/2" in diameter and has hours of clock time, a.m. and p.m., together with an arbitrary telescope time scale from 0 to 120 minutes. The fourth or inner dial remains fixed, after setting for your east-west position. The dial has half of the circumference divided into 12 parts representing the "hole number" settings on the telescope mount. A piece of 1/4" steel rod makes a good pivot for locating the dials centrally.

Scale and dial graduations can be laid out by a very simple method. Cut out a cardboard circle 29-1/16" in diameter from a packing box or other suitable material, and stretch a steel tape counterclockwise around the outer edge. (The circumference should be exactly 91 1/4".) Lay off months of the year using the following scale: 1/4" Jan. 1; 8" Feb. 1; 15" Mar. 1; 22 3/4" Apr. 1; 30 1/4" May 1; 38" June 1; 45 1/2" July 1; 53 3/4" Aug. 1; 61" Sept. 1; 68 1/2" Oct. 1; 76 1/4" Nov. 1; and 83 3/4" Dec. 1. Starting at January 1, lay off 31 days using 1/4" for each day, and number the days 0, 5, 10, 15, 20, 25, 30 in clockwise direction from February 1 to January 1. This protractor is used to lay out the days and months on the star and planet finder.

A second protractor 28 5/8" in diameter (90" circumference) will be required to lay out the other dials and scales. Stretch the steel tape around the outer edge in a counterclockwise direction and mark off 24 hours. Note that one hour equals 3 3/4", 2 equals 7 1/2", 3 equals 11 1/4", 4 equals 15". This protractor is used to lay out the right-ascension hour dials, the a.m.-p.m. clock time dial, and the telescope hole numbers. Next we mark off 120 minutes, in 5-minute steps (5 minutes equals 5/16") and lay out the telescope time scale. This minute protractor is also used to lay out the minutes of right ascension, minutes of clock time, and the finder telescope time scales. On the back side of the 28 5/8" disk lay off 10 10-degree steps at 2 1/2" intervals, and use this protractor to lay out the carriage declination scales. The declination index has 10 1-degree divisions; each degree on the circumference equals 1/4". The index should be laid out first, and the carriage 10-degree scale graduations and legends should be marked on the brown surface of the tempered



The star and planet finder, in five parts, enables rapid setting of the telescope for objects whose positions are given in the "American Ephemeris."

masonite using a good grade white crayon pencil (Eberhard Faber's extra thin Color-brite White #2111 is excellent for the purpose). Keep the point well sharpened in order to produce clean-cut lines. India ink will also make a good marking substance.

The setting of the star and planet finder is accomplished like the setting of a planisphere. First the day of the month is set, then the right ascension of the star or planet, then the clock time. All of these settings are then locked to the telescope mounting by means of one of the hourly telescope holes, and the slow-motion control takes over for as long as two hours. This device permits changing from one object to another with a minimum loss of time and telescope position. Alignment on the north pole of the sky may be only approximate, or may be accomplished as described in various books on observing and on telescope making (see page 162, **Making Your Own Telescope**, by Allyn J. Thompson).

Because of the long focal lengths used, eyepieces do not have to be very short and high powered. A 1 1/2", 1", and 3/4" should prove satisfactory for generally all work. These can be purchased in war-surplus catalogues, or separate lenses can be bought and mounted to make one's own eyepieces. For best results the eyepiece lenses should be of equal focal length and plano-convex, mounted with the convex sides facing each other and separated by about 7/10 of the focal length.

JAMES V. LAWRENCE
32-42 159th St.
Flushing 58, N. Y.

New Low Prices on PYREX Reflecting Telescope Kits

The most complete kits
on the market.

In addition to the usual supply of abrasives, rouge, etc., you get the new, fast-polishing cerium oxide to save hours of work.

You can get a brass diagonal holder (spider) for only \$1.00 additional if ordered with a telescope kit. Prices quoted below are for a Genuine Pyrex telescope blank and a plate glass tool.

PYREX MIRROR KITS

Diameter	Thickness	Price
4 1/4"	3/4"	\$ 5.50
6"	1"	\$ 8.00
8"	1 1/2"	\$11.00
10"	1 3/4"	\$19.00
12 1/2"	2 1/8"	\$35.50

PLATE GLASS KITS

Diameter	Thickness	Price
6"	1"	\$ 5.50
7"	1"	\$ 6.75
8"	1"	\$ 8.00

PRISM .. 6 1/2" long, 1 1/8" face .. \$3.25

PRISM .. 5 1/2" long, 1 1/2" face .. \$1.85

Postage Paid to 1st and 2nd postal zones from N. Y. Add 5% 3rd and 4th zones, 10% 5th and 6th zones. Add 15% 7th and 8th zones.

Send for free catalog of optical supplies.

DAVID WILLIAM WOLF

74 Hunnewell Ave. Elmont, L. I., N. Y.

OBSERVER'S PAGE

Universal time is used unless otherwise noted.

DEEP-SKY WONDERS

WHEN Admiral Smyth entered Xi Bootis on the pages of his famous *Celestial Cycle* (1844), there was much doubt about the true binary nature of the system. The three early observations (shown on the diagram as open circles), because of the error of the 1822 observation, suggested that the companion was moving in a straight line. Then the 1829 position indicated a very elongated ellipse. So the good admiral watched this pair with attention (his first and last observations are plotted here as crosses), and the

When the Aitken catalogue came out with measures to 1926, the true period obviously touched close to 150 years, the matter was settled, and the old observation of Herschel was justified.

Measures after 1926 on the diagram are from the card catalogue of Lick Observatory through the courtesy of Dr. H. M. Jeffers.

Xi Bootis is one of the few bright binaries that revolves fast enough so that amateur records will indicate changes over a decade or so. It lies at $14^h 49^m.1$, $+19^\circ 18'$ (1950); the components are of magnitude 4.7 and 6.6, spectra (by Mount Wilson) G5 and K5. If the amateur will fit his telescope with an eyepiece of high power which contains two parallel, slightly separated crosswires, and if this eyepiece be also equipped with an adjustable divided circle $3''$ to $5''$ in diameter carefully fitted so the center of the ocular field and the center of the circle coincide, then he can measure position angles easily and pretty accurately. His record book can then show this aspect of the binaries as he watches through the years. He will not need a clock drive or equatorial mounting, although these will help. All he needs do (but this must be done anew for each object examined) is to adjust his eyepiece until the stars trail across between the parallel wires. This gives the 270° - 90° zero mark for his circle. He then proceeds to measure the position angle of the double.

On the orbit diagram, the observations from 1866 to 1877 were all made by Dembowski who, without clock or micrometer, managed to make amazingly exact measurements which still are valuable today. So the amateur need not be dismayed by the elaborate equipment of the big observatories. In fact, the double-image micrometer described in *Amateur Telescope Making Advanced* can be constructed by anyone who has made a telescope, and it will give accurate results.


WALTER SCOTT HOUSTON

HINTS ON CHART CARE

The care of large sheets of paper, such as the Moon Sets, Skalnate Pleso *Atlas of the Heavens*, AAVSO charts, and the like, presents some difficulties that I have solved by binding the sheets between covers of tempered masonite.

This material is $\frac{1}{8}$ -inch thick, one side is smooth; it is strong and not too heavy, and sells for about 10 cents a square foot at lumber dealers. It is easy to saw with a fine-tooth handsaw or circular saw. For appearance, I have covered the boards with canvas, such as is used for folding chairs (obtainable in department stores), glued on. The bottom (back) cover is one piece, but the top (front) is cut lengthwise about two inches from its left edge. These two pieces are held together by a flexible hinge of Mystic tape (office-supply dealers) or glued linen. The covers and sheets are held together with aluminum ledger binding posts.

The Moon Sets were interleaved with tracing paper so that the formations could




The orbit of the visual binary star, Xi Bootis, from 1780 to the present. The 1780 and 1804 observations were by William Herschel.

accuracy of his work may be estimated from the way his plots fall on the curve. By the time he published his *Speculum Hartwellianum* (1860), he rejoiced that the binary nature of Xi Bootis was now beyond doubt. In another place he exclaims, "The physical connection of the components may therefore be deemed 'fully proven,' and that fact alone is a gratification to the contemplative mind."

The original William Herschel observation of 1780 is remarkably accurate when we consider the very amateurish nature of his instruments. He had no clock drive, no filar micrometer, not even an equatorial mounting—he used an altazimuth. Yet his position, checked against recent orbits, turns out to be only half a second of arc off in distance, and five degrees wrong in position angle. Using his position, the period comes out 152 years compared to the now accepted value of 151 years.

It was not until well into this century that the period was ascertained. Burnham's catalogue, with an orbit plot to 1905, remarks that it was still impossible to determine the period, and he gives references for orbits from 117 to 172 years, by various workers. Burnham, himself, thought the best observations indicated a period in the neighborhood of 175 years.



Fecker TELESCOPES

PRECISION BUILT
for the EXACTING AMATEUR
and the WORLD'S LARGEST
INSTALLATIONS

Designed and built for those
who want only the very best.
As distinguished in performance
as in its trim design. It's
everything you ever wanted in
a truly quality telescope.

PRICED
FROM
\$390.00

Write for free illustrated
catalog and prices, today!

FECKER
SINCE 1891

J. W. FECKER, INC.
2016 PERRYVILLE AVENUE
PITTSBURGH 12, PENNA.

TRIPLES 'SCOPE PERFORMANCE!

Sharper images at higher powers!
A startling statement positively
proven in 16-page telescopic educational
matter sent free on receipt of
self-addressed long envelope bearing
nine cents (9c) return postage.

First, the Goodwin Resolving Power lens
placed in front of eyepiece gives three
times the magnification on each by
lengthening your primary focal length
angle up to three times, yet extends eye-
piece out no more than two inches from
normal. This alone sharpens definitions.

Next, by achieving your highest powers
on more comfortable low-power eye-
pieces, you lessen image deteriorations
due to short-focus acute bending of the
convergent beam, since all usual eye-
pieces are $f/1$ or less.

Third, you get greater illumination and
wider field by relieving tiny aperture
restrictions of higher-power eyepieces.

The Resolving Power lens is achromatic,
coated, gives flat field sharp to the
edge. Here is astonishment in image
improvements! Price \$17.50 in 4" long chrome-plated
adapter tube fitting standard $1\frac{1}{4}$ "
eyepiece holders ONLY (but adaptable
to Unitrons). Money back if not
delighted after two weeks trial!

No COD's—Colleges and Observatories
may send purchase order.

FRANK GOODWIN

345 Belden Ave., Chicago 14, Ill.



BAUSCH & LOMB BALscope Sr.

Bausch & Lomb BALscope, Sr. is a highly precise, compact 60mm telescope. A deserved favorite for beginning astronomers, it also serves the advanced field as an adequate commercial version of a richest field telescope. Choice of four eyepiece powers as listed below. (With any of the three lower-power eyepieces, BALscope, Sr. is also an outstanding general observation terrestrial telescope.)

Eyepiece power	Exit Pupil diameter	Angular field
15×	4.0mm	2°40'
20×	3.05mm	2°5'
30×	2.0mm	1°28'
60×	1.0mm	0°33'

BALscope, Sr. with 15×, 20×, 30× or 60× eyepiece **\$95.00**

Extra eyepieces, each..... **25.00**

Tripod adapter (permits use of BALscope, Sr. with any pan-head camera tripod)... **6.85**

WRITE for descriptive folder. Bausch & Lomb Optical Co., 25818 Lomb Park, Rochester 2, N. Y.

be traced as observed and the name shown. An index was also made and bound in the front. The Skalnate Pleso plates are easier to handle if they are hinged with white Mystic tape; an extra strip of paper held in the binding allows the full plate to open easily and lie flat. Spring clips (such as Bulldog No. 2) will hold the sheets open against the front cover.

These bound sets are easily stored on a closet shelf, standing on the open side with the sheets hanging down (and thus kept straight) from their binding-post supports.

ROBERT J. BARR
2119 East Park Place
Milwaukee 11, Wis.

PREDICTIONS OF BRIGHT MINOR PLANET POSITIONS

Melete, 56, 9.7. June 12, 18:57.6 -7:35; 22, 18:51.6 -7:00. July 2, 18:44.2 -6:48; 12, 18:36.5 -6:59; 22, 18:29.9 -7:30. Aug. 1, 18:25.6 -8:19.

Hebe, 6, 8.2. June 22, 19:50.0 -7:19. July 2, 19:42.9 -8:09; 12, 19:34.1 -9:20; 22, 19:24.5 -10:50. Aug. 1, 19:15.4 -12:33; 11, 19:08.1 -14:19.

Melpomene, 18, 8.8. June 22, 19:58.4 -8:07. July 2, 19:51.9 -8:35; 12, 19:43.1 -9:25; 22, 19:33.3 -10:34. Aug. 1, 19:23.8 -11:58; 11, 19:16.0 -13:28.

Amphitrite, 29, 9.3. June 22, 20:29.7 -27:40. July 2, 20:22.9 -28:12; 12, 20:13.9 -28:41; 22, 20:03.7 -29:02. Aug. 1, 19:53.3 -29:12; 11, 19:44.1 -29:09.

After the asteroid's name are its number and the magnitude expected at opposition. At 10-day intervals are given its right ascension and declination (1958.0) for 0^h Universal time. In each case the motion of the asteroid is retrograde. Data supplied by the IAU Minor Planet Center at the University of Cincinnati Observatory.

SUNSPOT NUMBERS

March 1-2, 0, 0; 3, 1, 0; 4, 4, 0; 5, 0, 14; 6, 0, 0; 7, 0, 6; 8, 2, 0; 9-12, 0, 0; 13, 2, 7; 14, 7, 15; 15, 3, 8; 16, 0, 7; 17, 0, 0; 18, 3, 8; 19, 11, 10; 20, 11, 8; 21, 12, 10; 22, 11, 10; 23, 12, 10; 24, 12, 9; 25, 3, 9; 26, 0, 7; 27, 18, 17; 28, 21, 25; 29, 20, 32; 30, 48, 47; 31, 44, 48. Means for March: 7.9 American; 9.9 Zurich.

Daily values of the observed mean relative sunspot numbers are given above. The first are the American numbers computed by Neal J. Heines from Solar Division observations; the second are the Zurich Observatory numbers.

THE SUNSPOT MINIMUM

Minimum sunspot activity began late in 1952 when occasional zero values were in evidence. In 1953 January has yielded seven spotless days; February, 18 spotless days; and March, 11.

Prof. W. Gleissberg, director of the University Observatory, Bayazyt, Istanbul, Turkey, an authority on sunspot epochs, now predicts that minimum will occur during 1953, will be of short duration, and will be followed by a cycle slightly lower than the previous sunspot maximum of 1947, which was the second highest of all known sunspot cycles.

Up until November, 1950, although solar activity was declining, there had been no spotless days. At that time most minimum predictions were for late 1954 or 1955. The first spotless day of the current minimum occurred on December 19, 1950.

For comparison with this year's predicted minimum, the pattern of sunspot activity month by month during the pre-

vious minimum in 1944 is given here. The numbers of days with sunspots, month by month from January through December during 1944, were 10, 1, 13, 2, 4, 14, 11, 25, 24, 29, 22, 25. The numbers of days without spots for the same months were 21, 28, 18, 28, 27, 16, 20, 6, 6, 2, 8, 6.

NEAL J. HEINES
AAVSO Solar Division

MOON PHASES AND DISTANCE

Last quarter	June 4, 17:35
New moon	June 11, 14:55
First quarter	June 19, 12:01
Full moon	June 27, 3:29
Last quarter	July 3, 22:03

	June	Distance	Diameter
Perigee	5, 14 ^h	229,700 mi.	32' 20"
Apogee	18, 21 ^h	251,100 mi.	29' 34"

	July	Distance	Diameter
Perigee	1, 0 ^h	228,300 mi.	32' 32"

UNIVERSAL TIME (UT)

TIMES used on the Observer's Page are Greenwich civil or Universal time, unless otherwise noted. This is 24-hour time, from midnight to midnight; times greater than 12:00 are p.m. Subtract the following hours to convert to standard times in the United States: EST, 5; CST, 6; MST, 7; PST, 8. If necessary, add 24 hours to the UT before subtracting, and the result is your standard time on the day preceding the Greenwich date shown.

EVERYTHING FOR THE AMATEUR TELESCOPE MAKER

KITS \$4.50 and up

ALUMINIZING

Superior Reflecting Surface, Fine Finish. Will not Peel or Blister. Low Prices. Mirrors Tested | Prisms Eyepieces | Send for Free Accessories | Catalog

MIRRORS MADE TO ORDER

SEND FOR FREE PRICE LIST

Precision Optical Supply Co.

1001 E. 163rd St., New York 59, N. Y.

SKY-GAZERS EXCHANGE

Classified advertising costs 10 cents a word, including address; minimum charge \$2.50 per ad. Remittance must accompany order. Insertion is guaranteed only on copy received by the 20th of the second month before publication; otherwise, insertion will be made in next available issue. We cannot acknowledge classified ad orders. Sky Publishing Corporation assumes no responsibility for any statements made in the classified column, nor for the quality of merchandise advertised. Write Ad Dept., Sky and Telescope, Harvard Observatory, Cambridge 38, Mass.


FOR SALE: 10x Zeiss Teleplast binocular, extra stereo vision, \$55.00. 92-mm. terrestrial telescope, oculars 1 1/4" O.D., \$150.00. 3" scope, \$48.00. Norris Ingalls, 82 Purchase St., Newburyport, Mass.

NORTON'S "Star Atlas and Reference Handbook." Latest edition 1950, \$5.25. New publications: Gamow, "The Moon," \$2.75; Davidson, "Astronomy for Everyone," \$5.00. All foreign and domestic books. Write for list, Herbert A. Luft, 42-10 82nd St., Elmhurst 73, N. Y.

REFRACTORS: 4" Bardou, Unitron, Sans and Streiff. Complete with mounting and accessories that make observation a pleasure. Quality plus economy. Rasmussen and Reece, Amsterdam, N. Y.

6" f/8 REFLECTOR completely assembled in aluminum tube, finder. Write for snapshot, particulars; money back guarantee. Fred Messner, 107 N. Keeseey St., York, Pa.

FUTURE INQUIRIES pertaining to Hearn telescopes or The Telescope should be sent to A. H. Hearn Mfg. Co., 5924 N.W. 56 St., Oklahoma City, Okla. Prices and photo upon application. The finest 6" reflector you can buy at the price. Not sold through agents. A. H. Hearn, owner.



THE SUN, MOON, AND PLANETS THIS MONTH

The sun, on the ecliptic, is shown for the beginning and end of the month. The moon's symbols give its phase roughly, with the date marked alongside. Each planet is located for the middle of the month and for other dates shown.

Mercury, an evening object all month, attains greatest elongation on June 27th, $25^{\circ} 31'$ east of the sun. The elusive planet becomes visible to the unaided eye the first few days of June, setting an hour after the sun, at magnitude -1.0 . The best time to view Mercury will be in mid-June, when it will set over $1\frac{1}{2}$ hours after the sun, appearing at magnitude 0.

Venus attains greatest elongation in the morning sky on June 22nd, $45^{\circ} 46'$ west of the sun. At this time, the planet rises $2\frac{1}{2}$ hours before the sun, shining at magnitude -4.0 . Telescopically, Venus appears 50% illuminated, the disk $24''$ in diameter. The size of the disk is rapidly decreasing.

Earth arrives at heliocentric longitude 270° on June 21st at 17:00 UT. Summer commences in the Northern Hemisphere and winter in the Southern.

Mars will not be observable due to its proximity to the sun in the evening sky.

Jupiter reappears in the morning sky in the latter part of June, rising during the dawn.

Saturn, a fine evening object, will be on the meridian at 8 p.m. local time on June 11th. The planet will be located about $5\frac{1}{2}^{\circ}$ north of Spica in Virgo. Eastward motion will resume on June 24th, at which time Saturn will be at magnitude $+0.9$. The ring and satellite systems are always interesting telescopic objects. Minimum inclination of the rings occurs about June 18th, $11^{\circ} 9'$ to our line of sight.

Uranus will become invisible in the sun's glare as June progresses. On the 16th, Mercury will be in conjunction with Uranus, the latter $1^{\circ} 27'$ south.

Neptune may be observed until midnight, as an 8th-magnitude object approximately 1° south of Saturn in Virgo. The planet continues in retrograde motion and is at $13^h 20^m.9$, $-6^{\circ} 41'$ (1953) on the 15th.

E. O.

A JANUARY CONFIGURATION OF PLANETS AND THE MOON



The crescent moon, with earthshine on its darker side, Venus, and Mars are all in this photograph taken at the Forest Hall Observatory, Northumberland, England, on January 18th. The picture was transmitted by Frank J. Acfield. The camera, on loan from the British Astronomical Association, works at $f/5.8$, $19\frac{1}{2}$ -inch focal length, and was mounted on a 10-inch reflector. The exposure was 15 seconds on Ilford H.P.S. emulsion.

VARIABLE STAR MAXIMA

June 5, Z Ursae Majoris, 115158, 6.6; 9, S Hydrae, 084803, 7.9; 15, R Horologii, 025050, 6.0; 19, R Trianguli, 023133, 6.3; 23, S Coronae Borealis, 151731, 7.5; 25, T Centauri, 133633, 6.1; 29, R Andromedae, 001838, 7.0. July 2, R Lyncis, 065355, 7.9; 4, S Gruis, 221948, 7.8.

These predictions of variable star maxima are by the AAVSO. Only stars are included whose mean maximum magnitudes are brighter than magnitude 8.0. Some, but not all of them, are nearly as bright as maximum two or three weeks before and after the dates for maximum. The data given include, in order, the day of the month near which the maximum should occur, the star name, the star designation number, which gives the rough right ascension (first four figures) and declination (bold face if southern), and the predicted magnitude.

SKY-SCOPE

The new and improved $3\frac{1}{2}$ -inch Astronomical Telescope that amateurs everywhere are talking about.

Completely Assembled—\$29.75
Equatorially Mounted, 60 Power
 $\frac{1}{4}$ -wave Aluminized Mirror
Ramsden Type Ocular

125x & 35x Supp. Eyepieces, ea. \$5.00
6-power Finders postpaid, ea. \$7.30

We invite your attention to our free brochure describing in a straightforward manner the instrument's amazing performance.

THE SKYSCOPE CO., INC.

475-a Fifth Avenue, New York 17, N. Y.


Complete Telescopes and Supplies



Ramsden Eyepieces
Combination Eyepiece and Prism Holders
Mirror Cells
Finders
Equatorial Mountings
Prisms
Send for a price list

C. C. YOUNG

25 Richard Road, East Hartford 8, Conn.



The sky as seen from latitudes 20° to 40° south, at 9 p.m. and 8 p.m., local time, on the 7th and 23rd of September, respectively.

SOUTHERN STARS

FOR AN OBSERVER at 30° south latitude, at the times for which this chart is drawn, the brilliant portions of the southern Milky Way lie west of the meridian; the galactic center passed the zenith some 2½ hours earlier. Thus, a location in this latitude permits observation of the center of the Milky Way system under the best possible conditions,

and allows equally good conditions for study in all directions from the center.

As described by Jean Dufay in *Sky and Telescope*, December, 1952, page 41, the galactic center is located at about R.A. 17^h 30^m, Dec. -30°. For an observer at Philadelphia, this point cannot rise more than about 20 degrees above the southern horizon, and even for the 200-inch telescope this maximum distance is about 27 degrees.

In Arequipa, Peru, on the other hand,

where Harvard's Boyden station was first established in 1890, the galactic center passes only 14 degrees south of the zenith. At Bloemfontein, South Africa, where the Boyden station has been located since 1927, this distance is reduced to only one degree, probably less than the accuracy with which the direction to the center of the galaxy has been established. The Cordoba Observatory in Argentina is equally well situated, at latitude 31½° south.



STARS FOR JUNE

The sky as seen from latitudes 30° to 50° north, at 9 p.m. and 8 p.m., local time,

on the 7th and 23rd of June, respectively; also, at 7 p.m. and 6 p.m. on July 7th and 23rd. For other times, add or subtract ¼ hour per week. When fac-

ing north, hold "North" at the bottom; turn the chart correspondingly for other directions. The projection (stereographic) shows celestial co-ordinates as circles.



